



SOUTHERN RHODESIA.

REPORT

ON

The Public Health

FOR THE YEAR 1924.

Presented to the Legislative Assembly,
1925.

Salisbury, Rhodesia:
Printed by the Government Printer.

—
1925.

CONTENTS.

PART I.

CHAPTER I.

| | PAGE |
|--------------------------------|------|
| Administration | 1 |

CHAPTER II.

Work of the Department—

| | |
|--|----|
| Mines and Works Inspections | 5 |
| Health on Mines | 5 |
| Public Health Laboratory | 9 |
| Medical Inspection of Schools | 9 |
| Propaganda | 10 |
| Public Health Legislation | 10 |

CHAPTER III.

Vital Statistics—

| | |
|--|----|
| Marriages, Births and Deaths | 13 |
|--|----|

CHAPTER IV.

Public Health—

| | |
|--|----|
| Infectious, Communicable and Preventable Diseases ... | 14 |
| Notifiable Diseases, Malaria, Blackwater Fever, Pneumonia, Dysentery, Influenza, Small-pox, Trypanosomiasis, Enteric Fever, Malta Fever, Tuberculosis, Helminthic Diseases, Bilharzia, Tropical Ulcers, Venereal Diseases, Plague, Diseases peculiar to Natives. | |

CHAPTER V.

| | |
|---------------------------------------|----|
| Hospitals and Asylums | 21 |
|---------------------------------------|----|

PART II.

| | |
|--|----|
| Report on Pasteur Institute and Public Health Laboratory | 26 |
| Report on Ingutsheni Mental Hospital | 32 |
| Report on Administration of Habit-forming Drugs Regulations Proclamation | 34 |

PART III.

| | |
|----------------------------------|----|
| Tabular Appendix | 36 |
|----------------------------------|----|

The Reports of the District Medical Officers and Compound Inspectors have not been printed.



Digitized by the Internet Archive
in 2019 with funding from
Wellcome Library

<https://archive.org/details/b31477598>



SOUTHERN RHODESIA.

Report on the Public Health for the Year 1924.

Presented to the Legislative Assembly,
1925.

PART I.

CHAPTER I.—ADMINISTRATION.

The Public Health and Hospitals Department is in the Division and under the control of the Minister holding the portfolio of Colonial Secretary.

The *Authorised Establishment* consists of :—Medical Director, Government Bacteriologist, Medical Inspector of Schools, 5 Government Medical Officers (Grade 1), 4 Government Medical Officers (Grade 2), 15 Government Medical Officers (Grade 3), 1 aided Medical Officer, 1 chief clerk, 1 senior clerk, 1 Grade III. clerk, 5 lady clerks, 2 inspectors of compounds; senior matron, 7 matrons, 7 nurse-matrons, 29 qualified nurses, 25 probationers, 1 district nurse; Assistant Superintendent, Ingutsheni Mental Hospital; Assistant Superintendent, Morgenster Leper Settlement; 6 hospital secretaries and dispensers; 7 male attendants, Mental Hospital; 1 female attendant, Mental Hospital; 2 laboratory assistants.

Supplementary Auxiliary Staffs.—Six part-time hospital secretaries, 4 part-time secretaries to hospital advisory committees, 1 hospital orderly, 2 needlewomen.

The miscellaneous coloured and native staffs attached to the various institutions are as follows :—Four Indian cooks, 4 Indian laundrymen, 3 ward maids, 165 natives.

| | |
|--------------------------------------|-----|
| Total European Staff | 139 |
| Total Coloured Staff | 176 |
| <hr/> | |
| Total Staff | 315 |

It is with deep regret I have to record the death during the year of Dr. W. M. Eaton, O.B.E., the Assistant Medical Director. The late Dr. Eaton first joined the Government service in February, 1899, as District Surgeon, Hartley. Of late years he had held the post of

Assistant Medical Director, being stationed at Bulawayo, where he also held the appointment of Medical Superintendent of the Memorial Hospital and Medical Superintendent of the Ingutsheni Mental Hospital. Dr. Eaton had completed 25 years' service and would shortly have retired on pension. The late medical officer was a loyal and highly efficient servant of the Government, and the loss of such an able officer will be greatly deplored by all. The vacancy caused by Dr. Eaton's death has not yet been filled.

I also regret to record the death of Dr. S. Gurney, medical missionary, attached to the American Mission at Mrewa. Dr. Gurney was an aided medical officer, in that he was paid remuneration by the Government, in return for which he undertook all Government medical work in the Mtoko and Mrewa districts, and his services were also available for settlers under the special conditions laid down. The late Dr. Gurney was a man who could ill be spared. He at all times gave willing help to this Department in any direction where his medical knowledge and skill could be of use and was required, and his work as medical missionary at the various missions to which he was attached will long remain as a monument of life-long devotion to the welfare of suffering humanity. The American Mission authorities hope to be able to appoint a successor to the late Dr. Gurney at Mrewa, and when such appointment has eventuated it is hoped to renew the arrangement whereby such medical officer will be available for Government and public medical work at that centre.

During the year serious indisposition leave was granted to one medical officer, while vacation leave of absence was granted to four medical officers, one hospital secretary, one lady clerk, three Mental Hospital attendants and seven members of the nursing staff.

Dr. D. J. O'Keefe, Government medical officer, Mazoe, was retired on pension from 4th September, 1924, but he is continuing to hold his present post in a temporary capacity till the question of establishing a new Government station in a more central part of the Mazoe Valley has been finally decided. The payment of a pension to Dr. O'Keefe will remain in abeyance meanwhile. Dr. H. C. Titterton left the service in May, and Dr. Maurice Pearson was appointed to fill the vacancy.

The compound inspection staff still consists of three inspectors, though at the present time the staff has been depleted by one inspector having returned to duty in the Native Department in an emergency. It is intended to re-arrange these duties in the future so that two inspectors will suffice.

Eighty applications for appointment to the Southern Rhodesia nursing service were received from qualified nurses; nine were engaged. Five qualified nurses resigned during the year. Fifty-five applications were received for appointment as probationers, of whom thirteen were appointed. Four probationers resigned before completion of training, while an application has been received for an appointment three years hence, and a former member of the nursing staff now resident in Ceylon has written trying to arrange for her sister to be trained in the Salisbury Hospital. These appointments are now filled from the list of applicants resident in Southern Rhodesia.

Seven probationers successfully passed the Colonial Medical Council's examination for trained nurses during 1924. In this connection it is worthy of mention that since June, 1920, with one exception, probationers trained at the Salisbury Hospital have gained either a first or a second place in these examinations.

At the request of the London Tropical School of Medicine, I proceeded to England in September to discuss with the committee of the newly established Imperial Institute of Hygiene and Tropical Medicine the question of the establishment of a field branch of the London School in Southern Rhodesia. The outcome of the discussions which took place may be considered highly satisfactory. The Government of Southern Rhodesia will contribute £1,000 a year towards the scheme; it will also provide laboratory accommodation and equipment, and defray the cost of all materials in Rhodesia. The Government of Southern Rhodesia will also defray the cost of the journey of one research worker on one occasion to and from England, and the arrangement in the first instance will be for a period of three years. Workers sent by the school to the field branch will be given definite subjects and lines of research, which will be laid down for them by the director of the school in consultation with the Medical Director. A programme of work has been decided upon which will not be altered in any way unless with the joint agreement of the school authorities and this Department. The workers and the research work will in every particular be under the control of the director of the school, but the Medical Director will act as adviser to and be the local representative of the school. The research work to be undertaken in the first instance will be the continuation of Dr. J. G. Thomson's investigations into the causation and prevention of blackwater fever, and the question of the prevention of malaria in general, with which it is hoped to combine an entomological survey of the mosquitoes of the Colony. The appointment of Dr. Ross, Lecturer on Bacteriology at the Leeds University, as first Research Fellow has just been announced, and he is expected to arrive about the middle of April.

Much prominence has been given to this subject in the Press, and it is generally conceded that Southern Rhodesia is to be congratulated upon the fact that our capital has been chosen as the centre for the establishment of the first field station established by the London School of Hygiene and Tropical Medicine in any dominion of the Empire.

Financial.—The following figures show the expenditure under the Public Health and Hospital Votes under the respective headings for the year 1924, as compared with 1923 and 1922:—

Public Health (Expenditure).

| | 1924. £ | 1923. £ | 1922. £ |
|---|------------|------------|------------|
| Staff, salaries | 24,745 | 26,059 | 26,184 |
| Travelling expenses, Medical Director and staff, Government medical officers, Bacteriologist, compound inspectors, and rail and port charges | 3,673 | 3,694 | 3,832 |
| Treatment, maintenance and transport of lunatics, lepers and sick paupers, repression of infectious and contagious diseases, upkeep of lazarettos, purchase of vaccine lymph and quinine | 8,600 | 7,241 | 8,843 |

Public Health Laboratory and other Charges.

| | | | |
|-------------------------------------|---------|---------|---------|
| Expenditure | 400 | 329 | 395 |
| Administration of Foods and Drugs | | | |
| Ordinance | 191 | 256 | 261 |
| | <hr/> | | |
| | £37,609 | £37,579 | £39,515 |

Public Health (Revenue).

| | 1924. £ | 1923. £ | 1922. £ |
|--------------------------------------|------------|------------|------------|
| Bacteriological fees | 362 | 199 | 259 |
| Sales of quinine | 1,547 | 2,219 | 1,303 |
| | £1,909 | £2,418 | £1,562 |

Hospitals, Native Dispensaries and Asylums (Expenditure).

| | 1924. £ | 1923. £ | 1922. £ |
|---|------------|------------|------------|
| Salaries | 15,962 | 16,350 | 15,835 |
| Travelling expenses on appointment, duty or leave, rail and port charges | 1,440 | 855 | 1,308 |
| Provisions and medical comforts | 6,566 | 6,302 | 5,510 |
| Fuel, light and water | 2,892 | 2,856 | 2,694 |
| Furniture, equipment, repairs and clothing | 3,664 | 3,704 | 3,797 |
| Drugs, disinfectants and surgical appli- ances | 4,426 | 3,683 | 3,715 |
| Laundry | 1,130 | 1,040 | 605 |
| Sanitary fees | 455 | 482 | 348 |
| Produce, etc. | 7,764 | 8,101 | 9,207 |
| Miscellaneous | 387 | 514 | 554 |
| Grants-in-aid to hostels, hospitals and district nursing | 6,347 | 5,636 | 5,845 |
| | £51,033 | £49,523 | £49,418 |

Hospitals, Native Dispensaries and Asylums (Revenue).

| | 1924. £ | 1923. £ | 1922. £ |
|---|------------|------------|------------|
| Fees collected from paying patients | 13,326 | 15,055 | 13,812 |

The gross expenditure on Government hospitals, including Ingutsheni Mental Hospital, during 1924 amounted to £43,810, as compared with £43,229 in 1923, showing an increase of £581 on the previous year.

The revenue collected amounted to £13,326, as compared with £15,055 collected in 1923, showing a decrease of £1,729. This decrease is difficult to explain. The collection of hospital accounts is a matter which receives constant and careful attention, and I am satisfied there is little scope for improvement in the machinery of collecting revenue in this direction. The less satisfactory position may to some extent be due to a more stringent financial position generally during the period under review.

The gross earnings from paying patients amounted to £16,431, as compared with £16,603 for the previous year, this also showing a decrease of £172, while the total number of patients maintained was 5,594, as compared with 5,302 in 1923, an increase of 292.

In addition to the earnings from paying patients, the loss of revenue represented by the free treatment of paupers, police, gaol officials and others who were entitled to free medical attendance in Government hospitals, and for the cost of whom no inter-departmental charges are made, amounted to £10,011 15s., and if Ingutsheni Mental Hospital is included, this figure is increased to £19,142 10s., as compared with £16,141 for the previous year.

Complete statistics regarding the number of beds, daily average of patients treated, the revenue and expenditure, the average cost per patient per diem, the number of free patients, the number of units

treated, the cost to hospital votes of treatment and maintenance, worked out on the gross expenditure basis, and the loss of revenue represented thereby at the various institutions, will be found in Part III.

CHAPTER II.—WORK OF THE DEPARTMENT.

(1) *Mines and Works.*—(a) Inspections.—Towards the end of the year the compound inspector at Bulawayo was transferred back to the Native Department, and it has been decided to re-arrange the work so that if possible only two inspectors shall be required in future, with headquarters at Bulawayo and Salisbury respectively, thus bringing them into line with the inspectors of mines.

No report has been received from the inspector of compounds for Bulawayo for the period of the year in which he was acting, and I am, therefore, unable this year to show the total mileage covered or the mines visited. Inspections were, however, kept up till the advent of the heavy rains, which, owing to the impassable state of the roads, seriously curtailed their tours.

The reports of the compound inspectors on the whole indicated that the condition of native mine labourers was satisfactory. There were no prosecutions during the year under the Mines and Works Regulations or under the Native Labour Regulations, whilst complaints as regards non-payment of wages numbered only 18, as compared with 29 in 1923. There were six prosecutions with a like number of convictions on these counts; eight paid in full before prosecution, whilst four are being given time and are gradually paying off the arrears due.

These cases of non-payment of wages hardly fall within the scope of the duties of the compound inspectors, and are only undertaken by them as a convenience. They should by rights be dealt with directly by the Police, or the Native Commissioner of the district. One of the great difficulties in bringing these cases, and preventing the exploitation of the native for the benefit of the man who is gambling on what his prospect will produce, is to get the natives concerned to lodge a complaint, and the intervention of the compound inspector to explain their legal position to the natives has been helpful.

The diet regulations have been complied with in most instances to the best of the ability of the mines concerned. Owing to the abnormally low rainfall, however, difficulty has been experienced in many districts in supplying the ration of fresh vegetables which is required under the regulations, and the substitution of citrus fruits in lieu thereof has proved on the whole unsatisfactory, not on account of any shortage, but due wholly to faults in packing and costs of transport. These matters are now being taken up with the growers and the railway authorities respectively.

The inspector of compounds, Salisbury, has drawn attention to the fact that the amount of beans grown in the country is quite inadequate to supply the needs of the larger mines alone and that the prices ruling are altogether prohibitive, and it may be necessary to consider the withdrawal of this commodity from the compulsory scale of rations, which are fixed by regulation, and substitute some alternative.

(b) Health on Mines.—The number of employers rendering labour returns is as follows :—

| | 1923. | 1924. |
|----------------------------------|-------|-------|
| Mashonaland | 159 | 222 |
| Matabeleland | 174 | 218 |

The average size of the properties is indicated by the following table :—

| | |
|--|---------|
| Mines employing 2,000 natives and over | 3 |
| Mines employing 1,500 natives and over | 3 |
| Mines employing 1,000 natives and over | 4 |
| Mines employing 500 natives and over | 4 |
| Mines employing 400 natives and over | 1 |
| Mines employing 300 natives and over | 2 |
| Mines employing 200 natives and over | 11 |
| Mines employing 100 natives and over | 46 |
| Mines employing 50 natives and over | 71 |
| Mines employing 25 natives and over | 83 |
| Mines employing under 25 natives | 241 |
| Total | 469 |

The following table shows the number of cases of sickness, number of deaths, death rate per cent., sickness incidence per thousand per annum employed, and death rate per thousand per annum, amongst natives employed on mines in Southern Rhodesia for the year 1924 :—

Number employed, 41,286.

| Name of disease. | Total sick. | Total deaths. | Case mortality per centum. | Sickness incidence rate per mille per annum employed. | Death rate per mille per annum employed. |
|---------------------------------------|-------------|---------------|----------------------------|---|--|
| Malaria | 4,818 | 16 | .33 | 116.70 | .39 |
| Scurvy | 242 | 12 | 4.96 | 5.86 | .29 |
| Syphilis | 280 | 10 | 3.57 | 6.78 | .24 |
| Pneumonia | 1,791 | 330 | 18.43 | 43.38 | 7.97 |
| Phthisis | 101 | 24 | 23.76 | 2.45 | .58 |
| Other diseases of the chest | 2,445 | 39 | 1.59 | 59.22 | .94 |
| Dysentery | 205 | 15 | 7.32 | 4.97 | .36 |
| Diarrhœa | 406 | 2 | .49 | 9.83 | .05 |
| Other intestinal diseases | 282 | 25 | 8.87 | 6.83 | .66 |
| Heart disease | 57 | 31 | 54.39 | 1.38 | .75 |
| Debility | 344 | 14 | 4.07 | 8.33 | .34 |
| Influenza | 1,664 | 42 | 2.52 | 40.34 | 1.02 |
| Other diseases | 3,793 | 105 | 2.77 | 91.15 | 2.54 |
| Minor ailments | 8,689 | ... | ... | 210.59 | ... |
| Accidents | 460 | 89 | 19.35 | 11.12 | 2.16 |
| Totals | 25,577 | 754 | 2.95 | 619.51 | 18.26 |

The following table gives the mortality and causes of death amongst the various groups of natives employed on mines during 1924 :—

SOUTHERN RHODESIA.

MORTALITY ON MINES.

Annual Territorial Summary showing Mortality amongst Natives for year ended 31st December, 1924.

| Territorial classification Territories. | Average employed. | DEATHS. | | | | | | | | | | | | Death rate per mille per annum. | | | | | | | | | |
|--|-------------------|----------|---------|-----------|------------|-----------|------------------------------|----------------------------|------------|------------|-----------|------------|-----------------|---------------------------------|----------|--------|-------|-------|-------|-------|-------|-------|-------|
| | | Malaria. | Scurvy. | Syphilis. | Pneumonia. | Phthisis. | Other diseases of the chest. | Other intestinal diseases. | Dysentery. | Diarrhoea. | Debility. | Influenza. | Other diseases. | Accidents. | Disease. | Total. | 1924. | 1923. | 1924. | 1923. | 1924. | 1923. | |
| Southern Rhodesia | 11,133 | 5 | ... | 2 | 40 | 3 | 4 | ... | ... | 5 | 3 | 3 | 3 | 9 | 18 | 95 | 6.92 | 7.97 | 1.62 | 2.60 | 8.53 | 10.56 | |
| P.E. Africa | 4,069 | 1 | 1 | 1 | 19 | 7 | 3 | 3 | ... | 6 | 1 | 2 | 6 | 6 | 11 | 67 | 13.77 | 12.36 | 2.70 | 3.15 | 16.47 | 15.51 | |
| Northern Rhodesia | 10,967 | 5 | 5 | 3 | 96 | 9 | 17 | 6 | 1 | 7 | 9 | 1 | 11 | 39 | 17 | 226 | 19.06 | 15.06 | 1.55 | 2.27 | 20.61 | 18.17 | |
| Nyasaland | 14,649 | 5 | 6 | 4 | 173 | 5 | 15 | 6 | 1 | 7 | 17 | 8 | 22 | 46 | 41 | 356 | 21.50 | 16.20 | 2.80 | 3.01 | 24.30 | 19.21 | |
| Other sources | 554 | ... | ... | ... | ... | 2 | ... | ... | ... | ... | 1 | ... | 5 | 2 | 10 | 14.44 | 10.08 | 3.61 | 6.05 | 18.05 | 16.13 | | |
| Totals | ... | 41,372 | 16 | 12 | 10 | 330 | 24 | 39 | 15 | 2 | 25 | 31 | 14 | 42 | 105 | 89 | 754 | 16.07 | 13.29 | 2.15 | 2.77 | 18.22 | 16.05 |

Note.—The figure 41,372 is the average of the actual number of natives employed on the last day of each month, and not the daily average.

*Comparative Statement of Mortality amongst Natives Employed on Mines
in Southern Rhodesia, January to December, 1924.*

| Month. | Average No. of natives employed. | No. of deaths from disease. | Death rate per 1,000 per mensem from disease. | No. of deaths from accident. | Death rate per 1,000 per mensem from accident. | Total No. of deaths. | Death rate per 1,000 per mensem from all causes. |
|---------------|----------------------------------|-----------------------------|---|------------------------------|--|----------------------|--|
| January ... | 37,477 | 32 | .85 | 11 | .29 | 43 | 1.15 |
| February ... | 37,916 | 37 | .98 | 4 | .11 | 41 | 1.08 |
| March ... | 38,627 | 34 | .88 | 4 | .10 | 38 | .99 |
| April ... | 38,874 | 34 | .87 | 8 | .21 | 42 | 1.08 |
| May ... | 39,318 | 27 | .69 | 9 | .23 | 36 | .92 |
| June ... | 40,170 | 39 | .97 | 12 | .30 | 51 | 1.27 |
| July ... | 41,612 | 57 | 1.37 | 10 | .24 | 67 | 1.61 |
| August ... | 41,349 | 53 | 1.28 | 5 | .12 | 58 | 1.40 |
| September ... | 40,948 | 69 | 1.69 | 3 | .07 | 72 | 1.76 |
| October ... | 40,073 | 99 | 2.47 | 11 | .27 | 110 | 2.74 |
| November ... | 39,984 | 116 | 2.90 | 7 | .18 | 123 | 3.08 |
| December ... | 39,320 | 68 | 1.73 | 5 | .13 | 73 | 1.86 |


Totals and Averages.

| Year. | | | Per annum. | | Per annum. | | Per annum. |
|-------|-----|--------|------------|--------|------------|------|------------|
| 1924* | ... | 41,286 | 665 | 16.11 | 89 | 2.16 | 754 |
| 1923 | ... | 37,482 | 504 | 13.44 | 105 | 2.80 | 609 |
| 1922 | ... | 35,718 | 681 | 19.07 | 86 | 2.40 | 767 |
| 1921 | ... | 37,605 | 689 | 18.30 | 94 | 2.50 | 783 |
| 1920 | ... | 37,669 | 599 | 15.90 | 75 | 1.99 | 674 |
| 1919 | ... | 30,296 | 507 | 16.73 | 90 | 2.97 | 597 |
| 1918 | ... | 32,766 | 3,629 | 110.76 | 88 | 2.69 | 3,717 |
| 1917 | ... | 38,861 | 700 | 18.01 | 149 | 3.83 | 849 |
| 1916 | ... | 40,420 | 911 | 22.48 | 172 | 4.24 | 1,083 |
| 1915 | ... | 37,928 | 832 | 21.94 | 159 | 4.19 | 991 |
| 1914 | ... | 36,100 | 897 | 24.85 | 135 | 3.74 | 1,032 |
| 1913 | ... | 33,543 | 783 | 23.49 | 158 | 4.71 | 946 |
| 1912 | ... | 34,494 | 1,073 | 31.11 | 163 | 4.73 | 1,236 |
| 1911 | ... | 37,909 | 1,085 | 28.62 | 164 | 4.33 | 1,249 |

* Includes late returns.

MORTALITY ON MINES

Chart showing Death Rate per thousand per annum
among Native labourers from all diseases and from
Pneumonia and Influenza.



*Average Number Employed on Mines in Southern Rhodesia
during 1924, including late Returns.*

| | Natives. | Europeans. |
|--------------------------------------|------------|------------|
| Bulawayo | 12,681 | 449 |
| Gwelo | 9,528 | 445 |
| Victoria | 2,794 | 53 |
| Total for Matabeleland | 25,003 | 947 |
| Salisbury | 7,688 | 309 |
| Hartley | 6,772 | 287 |
| Umtali | 1,823 | 91 |
| Total for Mashonaland | 16,283 | 687 |
| Total for Southern Rhodesia | 41,286 | 1,634 |

The sickness and mortality rate amongst native mine labourers increased during the year, and it is interesting to note that the higher rate of sickness was ascribable to diseases which are associated with a short rainy season and a low rainfall, these being pneumonia and other diseases of the chest, scurvy, dysentery and other intestinal diseases.

The pneumonia rate was no doubt raised by a wave of epidemic influenza which appeared during the last quarter of the year and attacked natives employed on many of the larger mines, notably the Cam and Motor, Shamva, and Shabani Asbestos Mines. The mortality rate from diseases was raised in consequence, being 16.11 per thousand, as compared with 13.44 per thousand in 1923. On the other hand, the sickness rate from malaria was decreased. The average sickness and mortality rate still continues highest amongst labourers recruited from Nyasaland and the next highest amongst natives from Northern Rhodesia, and lowest amongst indigenous labour. It is somewhat significant, however, that amongst 41,372 native labourers employed on mines in Southern Rhodesia, only 11,133 are recruited in Southern Rhodesia.

(2) *Public Health Laboratory.*—The report of the Government Bacteriologist, which is to be found in Part II. of this report, shows a steady increase of the routine work of the laboratory.

Dr. J. G. Thomson, of the London Tropical School of Medicine, who was engaged in special research work into the aetiology of blackwater fever, returned to England in September, 1923, and it was not found possible to replace him. Following on the new arrangement that has been entered into between this Government and the London School of Hygiene and Tropical Medicine, the latter will in future keep one or more research workers in Rhodesia. Dr. G. R. Ross, M.B., Ch.B., D.P.H., Ph.D., Lecturer on Bacteriology at Leeds University, has been selected as the first Research Fellow.

During Dr. Orpen's absence on leave no research work was carried out in the laboratory, the staff in his absence being employed in routine work only. The gross earnings of the laboratory amounted to £862, as compared with £726 in 1923.

(3) *Medical Inspection of Schools.*—The Medical Inspector of Schools was transferred to this Department from the Education Department towards the end of the year, as it was considered that his work could be better correlated and developed under the Health Department. Dr. Gatchell's report, which has been separately printed and circulated, is

somewhat abbreviated this year on account of his absence on six months' leave. It, however, contains much food for thought, and should be carefully studied both by teachers and parents.

One of the most pressing needs of the school children is that of dental attention, which is receiving the serious consideration of the Government, and it is anticipated that provision will be made for this during the coming year.

(4) *Propaganda*.—Pamphlets prepared by the Department relating to malaria and blackwater, their causation and prevention, were in accordance with custom widely distributed, and arrangements were entered into with the Rhodesia Agricultural Union for a series of lectures by the district medical officers to farmers' associations on malaria and its prevention.

The following scientific papers were published by members of the staff :—

"Researches on Blackwater Fever in Southern Rhodesia," by J. G. Thomson, M.A., M.B., Ch.B., Director of Protozoology, School of Tropical Medicine, London, and Research Commissioner on Blackwater Fever in Southern Rhodesia, 1922 and 1923. This was published in London by the School of Tropical Medicine. It is a very complete history of Dr. Thomson's investigations in Rhodesia, and brings our knowledge of blackwater fever a big step forward.

"A Note on the Connection between Contagious Abortion and Undulant (Malta) Fever in Rhodesia," by L. J. J. Orpen, M.B., B.Ch., D.P.H. (Oxford), Government Bacteriologist, Southern Rhodesia. Published in the *South African Medical Record*.

"The Use of Saturated Organisms (Saturation Test) in Diagnosis," by L. J. J. Orpen, M.B., B.Ch., D.P.H. (Oxford), Government Bacteriologist, Southern Rhodesia. To be shortly published in the *South African Medical Record*.

"The Aëtiology of Blackwater Fever," by W. M. Hewetson, M.B., D.P.H., Government Medical Officer, Sinoia, Southern Rhodesia. Published in the *Journal of Tropical Medicine and Hygiene*.

"Some Observations on Blackwater Fever," by E. P. Carmody, M.B.E., M.R.C.S. (Eng.), L.R.C.P. (Lond.). Published in the *British Medical Journal*.

"Blackwater Fever," by O. E. Jackson, M.B., Bac. Surg. (Irel.). Published in the *South African Medical Record*.

"Chronic or Suppressed Malaria," by V. C. W. Vickers, M.R.C.S. (Eng.), L.R.C.P. (Lond.). Published in the *South African Medical Record*.

(5) *Public Health Legislation*.—(a) Public Health Act.—This Act, which consolidates and extends the public health legislation, was passed at the first session of the Legislative Assembly and became law on 1st January, 1925. Considerable discussion took place on the vaccination clauses, and amendments allowing for conscientious objectors were introduced, but apart from this the Bill passed the House in practically the form in which it was presented.

With the added powers now given to the Government and to local authorities, it is hoped that considerable strides will be made in matters which have long been awaiting attention. Regulations under this Act are now in course of preparation, and for a start will embrace vaccination, notification of infectious diseases, prevention of malaria and eradication of mosquitoes in towns, villages, mining camps and other communities, and the segregation and treatment of venereal diseases in natives.

(b) Southern Rhodesia Opium and Habit-forming Drugs Regulations.—A special report on the working of these regulations will be found in Part II. Certain clauses have been added to the regulations, allowing the keeping of limited stocks of opium by farmers and stock breeders for the treatment of sick animals, under permit issued and controlled by the Chief Veterinary Surgeon.

(c) Sale of Foods and Drugs Amendment Ordinance.—Regulations under this Ordinance, fixing standards for milk, butter and cheese, have been prepared, and are now under consideration, and it is anticipated that regulations dealing with these important commodities will be issued at an early date, as local authorities are pressing for their early introduction.

No whole-time Government analyst under the Foods and Drugs Act has yet been appointed, but it has been arranged, as a temporary measure, that this work will be undertaken by the chemical laboratory of the Agricultural Department.

(d) Natives Registration Ordinance, 1901, Amendment Ordinance, 1918.—This allows of the periodical medical examination of natives in employment in towns, but the regulations under this Ordinance still only apply to Salisbury, where, however, they have worked most satisfactorily, with undoubted benefit to the local authority concerned and to the public at large. Bulawayo is now adopting these regulations, and it is hoped that other municipalities will follow suit at an early date.

The following table, which has been supplied by the Medical Officer of Health for Salisbury, shows the number of natives examined and the diseases discovered and treated.

THE MUNICIPALITY OF SALISBURY.

Infectious Cases dealt with at the Lazaretto during 1924.

| Disease. | January. | February. | March. | April. | May. | June. | July. | August. | September. | October. | November. | December. | Total. | Case incidence per 1,000 per annum examined. | | |
|-----------------|----------|-----------|--------|--------|------|-------|-------|---------|------------|----------|-----------|-----------|--------|---|-----|-----|
| | | | | | | | | | | | | | | 5 | 6 | |
| Syphilis | ... | 10 | 8 | 7 | 5 | 3 | 5 | 3 | 5 | 6 | 5 | 5 | 72 | 5.93 | | |
| Gonorrhœa | ... | — | 3 | 1 | 2 | 1 | 2 | 2 | 1 | 1 | 1 | 1 | 17 | 1.40 | | |
| Leprosy | ... | — | — | — | — | — | — | — | — | — | — | — | 2 | .16 | | |
| Chicken-pox | ... | 2 | 1 | 2 | — | 1 | 1 | — | 1 | — | 1 | — | 9 | .74 | | |
| Itch | ... | ... | 12 | 10 | 10 | 8 | 2 | — | 2 | 9 | 2 | 5 | 4 | 6.01 | | |
| Filth | ... | ... | 1 | 1 | 2 | — | 1 | — | — | 1 | — | — | 6 | .49 | | |
| Chronic Ulcers | — | — | 1 | — | — | — | — | — | — | — | — | — | 1 | .08 | | |
| Boils | ... | — | — | 3 | — | — | — | — | — | — | — | — | 7 | .58 | | |
| Gonorrhœa Warts | — | — | 1 | — | — | — | — | — | — | — | — | — | 2 | .16 | | |
| Ringworm | ... | — | — | — | 1 | — | — | — | — | — | — | — | — | 2 | .16 | |
| Observation | ... | — | — | — | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | — | 2 | .58 | |
| Ulcers | ... | — | — | — | — | — | — | — | — | — | — | — | 1 | — | 2 | .16 |
| Measles | ... | — | — | — | — | — | — | — | — | — | — | — | — | — | — | .33 |
| Filth and Sores | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | .08 |

12

Total natives examined during the year, 12,136. Five operations were performed.

CHAPTER III.—VITAL STATISTICS.

The estimated European population at the end of June, 1924, was 37,519.

Marriages.—I am unable to make any reference to the marriage rate, as the registration of marriages is not dealt with by this Department.

Births.—There were 874 European births registered during the year, 446 being males and 428 females, giving a birth rate of 23.29 per cent., the lowest birth rate on record since 1920.

In addition the following births amongst Indians, coloured and natives were returned to the Registrar of Births :—

| | |
|----------|----|
| Indian | 54 |
| Coloured | 38 |
| Chinese | 1 |
| Native | 4 |
| | — |
| Total | 97 |

There were 15 plural births, of which 11 were European, two Indian, one coloured and one native.

The illegitimate births recorded numbered 33, being classified as follows :—

| | |
|----------|----|
| European | 21 |
| Coloured | 10 |
| Native | 2 |

In previous reports I have included a return showing the nationality of parents of the children born. This return is not included in this report, for the reason that the compilation of reliable statistics has been rendered difficult by the action of the informants, who either as nationalised British subjects or as residents of a British Colony now give their nationality as "British," instead of showing the country of their birth. Under these circumstances I have thought it desirable that this particular return should be discontinued.

Deaths.—Europeans.—There were 310 deaths recorded in 1924, as compared with 363 in 1923; of these, 188 were males and 122 females. The crude death rate was 8.26 per thousand of the population, which, corrected to the age and sex distribution in England and Wales, gives a corrected death rate of 10.03 per thousand. The mortality rate is therefore low, and compares favourably with the recorded death rates since the last census of the population, which was held in 1921.

| | | | | |
|------|------------------|-----------------|-------|--------------|
| 1921 | crude death rate | | 9.52 | per thousand |
| 1922 | crude death rate | | 8.98 | per thousand |
| 1923 | crude death rate | | 10.02 | per thousand |
| 1924 | crude death rate | | 8.26 | per thousand |

Infantile Mortality.—There were 65 deaths amongst children under one year of age, giving an infantile mortality rate of 74.49 per thousand births. This is a slight decrease from last year, but is still far too high. Fourteen deaths from broncho-pneumonia and pneumonia is considerably above the average, and may to some extent be explained by the wide-

spread epidemic of measles and whooping cough which occurred. Malaria as a cause of infantile mortality was, I am glad to say, reduced considerably, only three deaths from this cause being recorded, as compared with 12 in 1923.

The following table shows the causes of infantile mortality in the last four years :—

Causes of Death in Children under One Year of Age.

CHAPTER IV.—PUBLIC HEALTH.

Infectious, Communicable and Preventable Diseases.

Notifiable Diseases.—The following table summarises the weekly bulletins as issued by the Public Health Department:—

Table showing Infectious Diseases reported to the Public Health Department during 1924.

| Disease. | Number of cases. | | Number of deaths. | | Remarks. |
|---------------------------|------------------|---------|-------------------|---------|--|
| | European. | Native. | European. | Native. | |
| Chicken-pox | 103 | 388 | ... | ... | Many cases reported from Bulawayo and district; numbers not recorded |
| Cerebro-spinal meningitis | 1 | 3 | ... | 2 | |
| Diphtheria | 1 | ... | ... | ... | |
| Encephalitis lethargica | 1 | ... | ... | ... | |
| Enteric fever | 24 | 6 | 2 | 2 | |
| Erysipelas | 4 | ... | ... | ... | |
| German measles | 38 | 1 | ... | ... | |
| Influenza | 87 | 155 | 1 | 30 | Many cases reported from Lonely Mine district; numbers not recorded |
| Measles | 27 | 1 | ... | ... | Many cases reported from Zaka district; numbers not recorded |
| Mumps | 34 | 1 | ... | ... | |
| Puerperal septicæmia | 3 | ... | ... | ... | |
| Ringworm | 1 | ... | ... | ... | |
| Scarlet fever | 4 | ... | ... | ... | One suspected case reported from Umvuma |
| Scarlatina | 1 | ... | ... | ... | |
| Small-pox | ... | 2 | ... | ... | |
| Trypanosomiasis | 1 | 1 | ... | ... | |
| Whooping cough | 106 | 11 | ... | ... | |

Malaria.—The rise and fall in the incidence of malaria in relation to the character of the rainy season is well exemplified if we compare the year 1923 with that of 1924. In the former year we had a wet season, when the rains were above the average in the total amount of rain which fell, and late rains extended well into April and May. In 1924, on the other hand, the total rainfall was much below the average, and in the latter part of the rainy season it amounted to a drought in most districts. In 1923 we recorded a marked increase in the number of cases of malaria admitted to hospitals in the Colony, whilst in 1924 the admission rate on this account was much below the average, being 953 and 413 respectively.

The following table shows the admissions to all hospitals on account of malaria for the last four years, and the admission rate per thousand of the estimated population.

Malaria.

| Year. | Admissions to hospital. | Admission rate per 1,000 of the population. | Deaths registered. | Rainfall. | |
|-------|-------------------------|---|--------------------|-----------|----------|
| | | | | Season. | Average. |
| 1921 | 821 | 24.12 | 24 | 20.21 | 29.96 |
| 1922 | 500 | 14.35 | 12 | 21.22 | 16.11 |
| 1923 | 953 | 26.32 | 49 | 22.23 | 39.16 |
| 1924 | 413 | 11.11 | 13 | 23.24 | 16.69 |

Blackwater.

| | | | | | |
|------|----|------|----|-------|-------|
| 1921 | 53 | 1.58 | 23 | 20.21 | 29.96 |
| 1922 | 49 | 1.41 | 24 | 21.22 | 16.11 |
| 1923 | 63 | 1.74 | 36 | 22.23 | 39.16 |
| 1924 | 20 | 0.53 | 11 | 23.24 | 16.69 |

Though the malarial incidence showed such a marked decrease, I am afraid it can almost entirely be ascribed to the climatic conditions which existed, though no doubt individuals are more and more becoming alive to the necessity for the protection of themselves and their families. There has been no cessation in the educational and propaganda work which has now been for many years one of the chief instruments for combating its prevalence in rural areas. This year special lectures by the district medical officers to the various farmers' associations have been arranged and have apparently been much appreciated. The Public Health Act now allows of regulations being framed for the control of malaria and the eradication of mosquitoes, and regulations dealing with this have been prepared and will shortly come into force. These, however, can only in actual practice be applied to townships, villages, mining camps and other communities, and in the country districts the adoption of the simple precautions which are so necessary for his health must largely depend on the common sense of the individual. He at any rate cannot complain that he has not been fully advised.

Blackwater Fever.—The following table shows the admissions to hospitals and the mortality rate for the last 11 years on account of this serious sequel to malarial infection :—

| Year. | No. of cases of blackwater admitted to hospital. | No. of deaths in hospital. | Mortality rate per cent. |
|----------------|--|----------------------------|--------------------------|
| 1914 | 53 | 13 | 24.53 |
| 1915 | 62 | 16 | 25.81 |
| 1916 | 35 | 6 | 17.14 |
| 1917 | 48 | 13 | 27.08 |
| 1918 | 36 | 11 | 30.56 |
| 1919 | 37 | 7 | 18.92 |
| 1920 | 75 | 10 | 13.33 |
| 1921 | 53 | 6 | 11.32 |
| 1922 | 49 | 14 | 28.57 |
| 1923 | 64 | 14 | 21.88 |
| 1924 | 20 | 1 | 5.00 |
| | 532 | 111 | 20.86 |


As in malaria, the admissions to hospitals show a marked diminution for 1924, being only 20 cases with one death, as compared to 64 cases with 14 deaths in 1923.

The system of notification of blackwater fever to this office is now very complete, and in 1924 there were 39 cases notified with 11 deaths, giving a mortality rate of 28.2 per cent. In 1923 there were 122 cases notified with 40 deaths, with a mortality rate of 32.79 per cent. Here, as in the case of malaria, the short rainy season and period of drought in February, March and April were largely responsible for the diminution of blackwater fever in rural areas, and I am afraid that this satisfactory return cannot be accepted as a proof that this disease is in any way diminishing. The result of Dr. Thomson's researches has brought the study of the ætiology of blackwater fever a stage further, and it may now be accepted that in blackwater fever we are not dealing with a disease *sui generis*, but that it is a sequel of malaria, and generally, if not always, is associated with the particular form of malarial parasite known as the malignant tertian; but why haemolysis should occur in some cases of malaria and not in others, and what is the particular force which determines this cataclysm, are still unknown. Further research is required into this fatal and distressing disease, which is one of para-

Chart showing number of cases of Malaria and Blackwater Fever, with Rainfall, in Rhodesia during the years 1923 and 1924.

1923.

1924.



mount importance to the beneficial settlement of the more malarious parts of not only this Colony, but the whole continent of Africa, and the attention of the recently appointed branch of the London School of Tropical Medicine in Rhodesia will, in the first instance, be directed to this subject.

Quinine.—The importation of quinine by the Government and its distribution to the public through the agencies of post offices, stores, etc., has been continued throughout the year, and is now widely taken advantage of. Some 8,954 bottles of 100 tablets each, representing 895,400 5-grain tablets of quinine, were distributed by this Department during the year, and were sold to the public at 4s. 6d. per bottle.

The indigenous native is gradually becoming alive to the value of this drug, and the demand for quinine by natives is gradually increasing. In order to meet this, it is proposed to sell small tubes, each containing five 5-grain tablets of quinine, to the natives at 3d. a tube, and the Chief Native Commissioner anticipates that this will be widely taken advantage of.

Pneumonia.—There was a general increase in the incidence and mortality rate from pneumonia both amongst Europeans and natives, and especially affecting natives employed in mining. In many instances this was secondary to influenzal infection, being specially marked in the last quarter of the year. It is a noticeable fact, however, that whilst the incidence of malaria and blackwater increases in seasons of high rainfall, pneumonia, dysentery and scurvy tend to be highest in years following low rainfalls, and this year several of the Government medical officers in their annual reports have drawn attention to the abnormal amount of pneumonia this year.

Dysentery.—As in previous years of drought or low rainfall, there was an increase in dysenteries and diarrhoeal diseases admitted to hospitals. In opposition to pneumonia, which in this Colony is essentially an industrial disease, dysentery affects the native in his kraal more than in service, children being more prone to infection than adults. In the majority of the cases seen it was bacillary in type.

Influenza.—Following on the great epidemic of influenza in 1918, this country, as in England, has been subjected to mild recurrent epidemics which affect Europeans and natives alike, but the mortality is higher amongst the latter, owing to secondary pneumonia. These waves of recurring influenza were more prevalent in the latter part of the year, that is, towards the end of the dry season. There were 91 Europeans and 332 natives admitted to hospitals on this account during the year, with 30 native and no European deaths. In 1923 the admissions on this account were 63 and 131 respectively. Amongst natives employed on mines there were 1,664 cases recorded, with 42 deaths, as compared with 1,341 cases in 1923, with 29 deaths.

Small-pox.—Two cases of small-pox were recorded during the year, both in natives. This is the lowest return on record. There were no deaths.

Vaccination.—36,193 vaccinations amongst the native population were carried out last year. Owing to the prevalence of small-pox in the Portuguese territories adjacent to our borders, a system has been inaugurated this year of vaccinating at the port of entry all alien natives entering this Colony in search of work, 16,552 of these natives being vaccinated immediately on entering the country during last year. This should help to limit the number and extent of the epidemics of this

disease, which in the past have been of almost annual occurrence, and which have frequently been traced to outbreaks in bordering countries.

The average of unvaccinated children attending schools is reported by the Medical Inspector of Schools to be 17.11 per cent. of the children examined. A large number of these are the children of parents who do not realise the importance of this in early childhood or who are resident in country districts far from a doctor. Arrangements are now made for children to be vaccinated either by the Medical Inspector of Schools when on his visit of inspection or in town schools by the schools medical officer, and in the districts by the district medical officer, who annually travels round the schools of his district for the purpose.

The clauses in the Public Health Act dealing with vaccination came in for a considerable amount of discussion when this Bill was before the Legislative Assembly, and resulted in the Government introducing additional clauses providing for the conscientious objector. It is not, however, anticipated that this will be widely taken advantage of.

Trypanosomiasis.—Two cases of sleeping sickness were reported, one a European and one a native. The European was on a shooting trip ostensibly in the Mafungabusi area, but in spite of being warned he penetrated into the Sebungwe area in the neighbourhood of the Sipoli Vlei, where several cases of sleeping sickness have occurred in previous years. He became infected and was removed to Bulawayo, where he was treated with Bayer 205 and discharged ostensibly cured. He has since left the Colony, and up to now no further record of his state of health has been procurable. The native case was working in the Mafungabusi area, and prior to that came from Loangwa Valley, Northern Rhodesia, and the source of his infection is obscure. The Mafungabusi native district has been under suspicion for some time, and Dr. A. J. Mackenzie, the senior Government medical officer at Gatooma, carried out an exhaustive examination of all the natives in this area, but with negative results, and it is just possible that this was a latent case, infection having been contracted prior to entry into Southern Rhodesia. He was sent to Salisbury and given a course of Bayer 205, which benefited him at first to the extent that he appeared to all intents and purposes cured, but his symptoms somewhat suddenly relapsed, and he eventually died of the disease early in 1925.

Enteric Fever.—This disease shows a gratifying tendency towards decrease, and in spite of the fact that the sanitary services of most of our towns are of a somewhat primitive character, typhoid fever cannot be said to be a disease of outstanding importance, whilst it is noticeably absent in the native races, except those which are brought into close contact with European civilisation.

The admissions to hospital and deaths from this cause for the last five years are as follows :—

| Year. | Admissions. | Deaths. |
|----------------|-------------|---------|
| 1920 | 74 | 14 |
| 1921 | 53 | 9 |
| 1922 | 46 | 4 |
| 1923 | 61 | 11 |
| 1924 | 42 | 7 |
| | — | — |
| | 276 | 45 |

Malta Fever.—This not being a notifiable disease, we have no record of the actual number of cases occurring, but admissions to hospitals continue to increase, there being 16 admissions in 1924, as compared

with 12 in 1923. It is noticeable that natives appear to be practically immune, no case in a native having yet been recorded.

Research into the causal agent and its relation to contagious abortion in cattle is being continued, but was suspended during Dr. Orpen's absence on leave.

Tuberculosis.—This malady tends to show a slight but steady increase, the following table giving the deaths registered from all forms of tuberculosis in the last eleven years :—

| Year. | No. of deaths. |
|-------|----------------|
| 1914 | 57 |
| 1915 | 58 |
| 1916 | 52 |
| 1917 | 74 |
| 1918 | 83 |
| 1919 | 88 |
| 1920 | 92 |
| 1921 | 103 |
| 1922 | 87 |
| 1923 | 73 |
| 1924 | 94 |
| | 861 |

There were 152 admissions to hospitals in 1924, as compared with 142 in 1923, of which 20 were Europeans and 132 natives.

The deaths were 94, as compared with 73 in 1923 and 87 in 1922. They were classified as follows :—

| Disease. | 1922. | 1923. | 1924. |
|----------------------------|-------|-------|-------|
| Tuberculosis of the lungs | 73 | 63 | 85 |
| Tubercular meningitis | 4 | 3 | 1 |
| Abdominal tuberculosis | 2 | 5 | 2 |
| Pott's disease | 3 | ... | 2 |
| Tuberculosis, other organs | 4 | ... | 1 |
| Disseminated tuberculosis | 1 | ... | 2 |
| White swellings | ... | 1 | ... |
| Acute miliary tuberculosis | ... | 1 | 1 |

The increase in phthisis is largely confined to native male adults in employment, and so far as Europeans are concerned shows a tendency to decrease, this being assisted by the strict surveillance of immigrants and the exclusion of persons suffering from this disease in an active form.

Helminthic Diseases.—*Bilharzia.*—The Government Bacteriologist reports 63 examinations of urine for this disease, with 22 positives. Some research amongst the snails implicated as vectors has also been done, and results will be found in the report of the Public Health Laboratory.

The Medical Inspector of Schools reports 17 cases amongst the children he examined last year, which is over and above the 71 cases previously discovered by him. He rightly draws attention to the carelessness of parents in this respect. Bilharziasis is a serious disease, affecting the general health and nutrition of the child and seriously handicapping the prospects of life, and cannot be too carefully avoided. When discovered, the disease should be treated in the earliest stages.

The following table shows the admissions to hospitals on account of helminthic diseases :—

| Disease. | Europeans. | Natives. | Totals. |
|--------------------------|------------|----------|---------|
| Tæmia solium | 4 | 2 | 6 |
| Tæmia saginata | ... | 2 | 2 |
| Hydatids | ... | 1 | 1 |
| Bilharzia | 10 | 10 | 20 |

Tropical Ulcers.—The occurrence of tropical ulcers in natives shows no tendency to decrease, and adds appreciably to the sickness rate on mines, besides being a source of considerable economic loss to the industry. Dr. F. H. Ellis, in his report on the Salisbury Hospital, draws attention to the successful treatment of these ulcers by the application of the leaves of the stramonium plant, which is a common weed on the veld.

Venereal Disease.—Government medical officers, Native Commissioners and missionaries continually report the prevalence of venereal disease amongst the native population, this being more extensive in some districts than in others, but unfortunately we are not able at present to collect sufficiently reliable figures which might indicate whether venereal diseases are increasing or not amongst the native population. On the mines the recorded cases show a slight increase in relation to the number employed, being .67 per cent. in 1924, as compared with .59 per cent. in 1923; but, apart from this, there is no reason to believe that the ratio is markedly altering. Provision for the segregation and treatment of these cases is, however, urgently required, and the additional powers now given to the Government and local authorities under the recently promulgated Public Health Act will materially assist in allowing for this. Certain proposals regarding the establishment of venereal clinics in selected areas are, in fact, now receiving the consideration of the Government.

Plague.—Though bubonic plague has so far not appeared in Southern Rhodesia, its continued spread through the agency of veld rodents in the Union of South Africa cannot but be a source of considerable anxiety. Should the epidemic amongst the wild rodents continue to spread, it is difficult to see how we are to effectively protect our own borders.

In July of last year I visited the plague infested area of the Free State and Transvaal for the express purpose of studying the conditions prevailing and consulting with the Union Public Health authorities, and the result of these enquiries was embodied in a memorandum which was submitted to Ministers.

Every precaution against the introduction of plague, either by persons or rodents, by rail, is being taken, and recommendations for the reduction of the rat population in towns and villages were published for general information.

Diseases Peculiar to Natives.—In the annual report for last year I drew attention to certain diseases of unknown origin which were peculiar to natives and which required further investigation, and I specially mentioned splenic abscesses, endemic cirrhosis of the liver and chronic lymphangitis of the lower extremities; whilst Dr. Ellis reports this year a case of onyalia in a native—a rare disease of unknown origin, which up to now has only been reported from Portuguese West Africa.

I regret that it has not yet been possible to institute any enquiries into these diseases, but hope that steps will be taken in the near future to extend our knowledge of the ætiology of these diseases, especially with regard to the condition of chronic lymphangitis of the legs, which is extremely common and is the cause of a large amount of invalidism amongst our native employees on mines and other works.

CHAPTER V.—HOSPITALS AND ASYLUMS.

There were 6,824 admissions to general hospitals in Southern Rhodesia in 1924, as compared with 6,325 in 1923. Of these, 2,833 were Europeans and 3,991 were coloured and natives. European admissions decreased by 477, while native and coloured admissions increased by 976.

European admissions reached their highest in the month of May and their lowest in the month of July, being approximately similar to observations made in former years, while native admissions were highest in October and November and lowest in July.

Memorial Hospital, Bulawayo.—Negotiations bearing on the future of this institution have been in progress for some time, and it has now been finally decided that as from the 1st of April, 1925, the hospital will be taken over by the Government, which will from that date be responsible for its administration and upkeep. The hospital board will continue, but as an advisory committee only, in accordance with the practice at other hospitals. Fees charged to patients will be reduced to the Government scale, and the medical superintendent will be a Government medical officer, who will reside on the premises, and the institution will be absorbed in the hospital system of the Colony.

The total number of white patients admitted to the Memorial Hospital during the year was 731, as compared with 925 in 1923, and 1,003 natives, as compared with 682 in 1923. Three hundred and forty-seven European (174 major and 173 minor) and 82 native (32 major and 50 minor) operations were performed during the year.

Salisbury Hospital.—There were 2,122 cases admitted during the year, of whom 1,093 were Europeans and 1,029 were natives and coloured, as compared with 1,072 Europeans and 808 coloured and natives in 1923. The figure 2,122 constitutes a record, and shows a substantial increase in the work done at this institution. The operations performed were 553 European (201 major and 352 minor) and 97 native (28 major and 69 minor).

The only constructional work done during the year was the erection of an isolation block. The provision of accommodation for cases of infectious disease developing after a patient's admission to hospital has been long required, and the building in question has already proved its usefulness. The water supply of the hospital has further been supplemented by the erection of an electric pump at the borehole well, which is working satisfactorily, and it is anticipated will effect a substantial saving in expenditure.

Additional accommodation for Europeans is becoming a matter of urgency, the congestion in the women's ward being really serious at times, whilst it is very desirable that further separate accommodation for children should be provided. The erection of an additional wing to this building would meet all immediate requirements, and it is hoped that this extension will be provided in the near future.

On two occasions during the year articles appeared in a Salisbury weekly paper attacking the hospital and its management. The subject matter of the articles in question received very full and impartial investigation, and the result of the enquiries elicited the information that the complaints stated to exist were not well founded, and the hospital advisory committee, who were also asked to enquire into the matter, expressed the opinion that the Salisbury Hospital is an exceedingly well-arranged institution, that the members have sought opinions from patients and made efforts to find any real source of dissatisfaction with-

out result. The committee have particularly asked that this expression of satisfaction with the management, as evidenced by the committee's own investigations and the opinions of patients, be embodied in this report.

Umtali Hospital.—There were 593 admissions, of whom 308 were Europeans and 285 were natives, as against 463 Europeans and 137 natives for the previous year. The senior Government medical officer, Umtali, remarks that the decrease in the number of Europeans was due to the smaller number of malarial cases which followed the short rainfall, and that the increase in the number of native patients was due to the greater number seeking admission owing to a recognition of the benefit of European treatment in hospitals. Twenty-eight operations were performed, 20 being Europeans and 8 natives.

Certain structural alterations were undertaken, whereby the old nurses' home was converted into wards for women, and the house formerly occupied by the doctor was altered and added to to form a nurses' home. This re-arrangement has temporarily relieved the pressing need of more accommodation for female patients, but it cannot be denied that the present hospital buildings are no longer suited to present-day requirements, and their continued use can only be justified till funds can be provided for the erection of a new hospital suited to the needs and importance of the situation. Increased accommodation for natives is also urgently required, but it would be unwise to incur the expense of adding to the present buildings.

It has been suggested that the present hospital buildings might continue to be utilised as a hospital for Indians, coloured and natives, a portion of the existing European block being set aside for the former, but the establishment of two separate hospitals would entail a higher expenditure than if both hospitals were administered as one institution in the same grounds, and I am of opinion that the advantages to be gained thereby would counterbalance the initial expenditure which would be saved if it were decided to continue to use the present hospital buildings for natives when the new hospital for Europeans is erected. A site for a new hospital has been provisionally selected.

Gwelo Hospital.—Seven hundred and thirty-five patients were admitted during the year, of whom 167 were Europeans and 565 were natives, as against 218 Europeans and 400 natives for the previous year. The operations performed numbered 38 European (16 major and 22 minor) and 128 native (19 major and 109 minor).

The number of natives treated in this hospital is in excess of the accommodation available, and additional native ward accommodation is an urgent necessity. The sanitary system of this hospital is also unsatisfactory. These matters are, however, receiving close attention, and it is hoped that the work of erecting additional native wards, as the nucleus of a new native hospital to be erected in the grounds of the present hospital buildings, will be put in hand at an early date. It is also hoped that an improved and more up-to-date sanitary system will be devised in the near future.

Fort Victoria Hospital.—The admissions were much the same as in the previous year, there being 104 Europeans and 109 natives, as compared with 98 Europeans and 110 natives in 1923. European operations numbered 14 (6 major and 8 minor), and operations in native cases totalled 4 (3 major and 1 minor). The Government medical officer, Fort Victoria, also reports that there were 764 out-patient cases, many natives

attending for the dressing of injuries received, usually in beer drinks. The work of building the new hospital at Fort Victoria has not yet been commenced, but it is expected this work will be put in hand early in the new year.

Gwanda Hospital.—The admissions to this hospital numbered 42 whites and 156 natives, as compared with 36 Europeans and 124 natives in 1923. Three European (minor) and 9 native (3 major and 6 minor) operations were performed.

Enkeldoorn Hospital.—Fifty-two patients were admitted during the year, of whom 13 were white and 39 were natives, as compared with 32 whites and 24 natives for the previous year. Nine minor operations (5 white and 4 native) were performed.

Gatooma Hospital.—It has been decided by the Government that a new general hospital be erected at Gatooma, and this work will probably be commenced early in the new year. A site has been selected bordering the township, and the European and native hospitals will exist as one institution. The removal of the native hospital from the centre of the township will dispose of a constant source of grievance on the part of the townspeople, and the establishment of both hospitals as one institution will tend to economy in administration and to efficiency. No accommodation for maternity cases will be provided in the new hospital, and this has led the inhabitants of Gatooma to advance a plea that accommodation for such cases be provided in the new hospital when it is erected. The local committee has been informed, however, that the policy of the Government in the past has been to assist local effort in maintaining maternity homes by the payment of a substantial grant-in-aid, and it is difficult to find a reason why Gatooma should be on a different footing to Salisbury, Gwelo, Umtali, Fort Victoria, Hartley and Selukwe in this respect. During the year 222 patients were admitted to the European hospital and 472 patients were admitted to the native hospital, as compared with 262 Europeans and 414 natives for the previous year. No operations were performed on Europeans, while 13 major and 56 minor operations were performed on natives. In the maternity wards attached to the present hospital there were 19 confinements during the year.

Shamva Hospital.—Eighty white patients were admitted and 176 natives, as compared with 125 whites and 202 natives for the previous year. Fifteen minor operations were performed on Europeans and 6 major and 25 minor operations were performed on natives. The new hospital is not yet erected, but it is expected that this work will be put in hand at an early date.

Sinoia Hospital.—It has been decided to erect a new hospital at Sinoia, and this work will probably commence early in the new year. The present hospital buildings will remain and will be used as the nurses' home. The white admissions numbered 67 and the native admissions 103, as against 73 and 82 respectively for the previous year. The smaller number of European patients is attributed to a falling off in the number of admissions on account of malaria. Two operations on Europeans were performed. Five cases were admitted to the maternity ward.

Belingwe Hospital.—This small cottage hospital, which is under the care of the Government medical officer, Filabusi, who visits at regular intervals, continues to supply a want of that district. Sixty patients were admitted during the year, of whom 6 were Europeans and 54 were natives, as compared with 6 and 32 respectively for the previous year.

The residents of Shabani have asked that a Government hospital be erected at Shabani. It is obvious that, with a Government hospital established at Belingwe, there would be no justification for a second Government hospital in the same district, only 20 miles distant, and that, if the Government erected a hospital at Shabani, the institution at Belingwe should be closed. Urgent reasons have, however, been advanced why the Belingwe hospital should not be moved, and I am inclined to agree with this view. It is to be borne in mind that the populace at Shabani consists for the most part of persons employed by the groups of mines working there, and on the mines concerned rests the responsibility of providing adequate arrangements for the nursing and treatment of their employees when sick. This view has been communicated to the people concerned, who have been informed that, if they will combine and erect a hospital at Shabani, the Government will assist by contributing a grant-in-aid, and there the matter rests for the time being.

Morgenster Leper Settlement.—The number of lepers at the settlement as at the 31st December, 1923, was 128 (86 males and 42 females), 124 being adults and 4 being children. During 1924 there were 78 new cases admitted (55 males and 23 females). One male was discharged, 9 escaped (8 males and 1 female), and 4 were allowed out on probation (1 male and 3 females). Fifteen lepers died, of whom 8 were males and 7 were females. The daily average treated numbered approximately 55, and the daily average maintained was 165. The number of lepers in the settlement on the 31st December, 1924, was 177 (123 males and 54 females). The superintendent's report says that the behaviour of the lepers has been good.

Under the provisions of the Leprosy Repression Ordinance the leper asylum is visited at least once in each year by a board of officers appointed by the Government. The board reported that when they inspected the settlement conditions were satisfactory, but recommended that certain of the lepers who were in need of constant medical supervision should be transferred to the Ndanga native dispensary, where they suggested separate ward accommodation for such cases should be provided. The board also suggested that arrangements should be made for the removal and care of children born of leprous parents. It has not yet been possible to do anything in regard to the transfer of lepers to Ndanga, and the latter suggestion is receiving attention. The present site of the leper settlement is not altogether suitable, and it has been decided to remove the settlement to a new site, and an area, being part of what is known as the Nyanda block, situated some 14 miles south-east of Fort Victoria, has been selected. This site will be more convenient in every respect, besides allowing ample room for expansion, and has the great advantage of being outside the native reserve and not hedged in with kraals. Survey and building operations will take time, but it is hoped that the removal of the settlement to its new home will be accomplished before the end of next year.

Ingutsheni Mental Hospital.—In his report for the year, which will be found printed in Part II. of this report, the medical superintendent again draws attention to the overcrowding and the urgent need of additional accommodation. Accommodation also is still required for European female patients. In this connection I can only endorse the views expressed by the medical superintendent, and hope that the necessary funds to provide for these real and urgent requirements will be forthcoming, so that the work of providing the necessary buildings may not be longer delayed.

The superintendent has for some time past urged the necessity of employing only trained mental nurses as attendants. The assistant superintendent and the two senior attendants are qualified men; two further certificated men are shortly to be engaged in England, and it is intended that the unqualified men at present employed shall gradually be replaced by trained men, so that the whole of the nursing staff shall consist solely of attendants qualified by training and examination in the nursing and care of the insane.

Statistics relating to the principal diseases treated in Government hospitals during the year will be found in the appendix to this report.

A. M. FLEMING,
Medical Director.

PART II.**Pasteur Institute and Public Health Laboratory.****REPORT FOR THE YEAR 1924.**

Staff.—The permanent staff consists of the Bacteriologist, senior assistant, junior assistant, and two native laboratory boys, while it is hoped that an Analytical Chemist and perhaps an Entomologist will also be appointed. For purposes of research special investigators are sent out from the London School of Tropical Medicine, and this year it is hoped that a Bio-Chemist and an Entomologist will be sent out. The services of Dr. Ross have been secured so far for 1925.

Pasteur Institute.

The fresh strain of rabbits proved quite satisfactory, and there has been no sickness among them or the guinea-pigs.

One patient (coloured) from Portuguese East Africa was treated here; a course of treatment was sent to a patient in Nyasaland, and five courses of treatment were sent to Livingstone. We have had no patients from Southern Rhodesia since 1913.

Our new methods, both of preparing and of sending out virus for treatment, have proved very satisfactory in every way. The virus is kept as a glycerine emulsion which can be used for passage inoculations or for treatment at any moment, while courses of treatment, ready for use, are posted to doctors whenever required, so that patients can be treated at their own homes instead of having to come to Salisbury for several weeks.

Public Health Laboratory.**I.—ANALYSIS OF WORK DONE.**

A. Research.—No research investigator could be sent out from the London School of Tropical Medicine last year to continue the research work on blackwater fever. Dr. Ross is expected to arrive in March of this year to carry on this work.

Owing to the absence of the senior assistant during the first part of the year and of the Bacteriologist during the second half, no continuous research could be maintained by ourselves, but such work as was possible was done on malaria, bilharzia and the diagnosis of Malta and enteric-like fevers, which will be described later under these headings.

B. Routine.—The work increases steadily, 1,714 examinations having been made in 1924, as opposed to 1,457 in 1923 and 807 in 1922.

Outside districts are making increased use of the laboratory, as is shown in the following table:—

| | 1923. | 1924. |
|------------------------------------|-------|-----------------|
| Southern Rhodesia— | | |
| Salisbury | 1,352 | 1,487 |
| Nineteen other towns | 99 | 21 towns 227 |
| Northern Rhodesia | 3 | ... |
| Portuguese East Africa | 3 | ... |

These figures for places outside Salisbury, though improved, are not yet satisfactory. Many specimens cannot travel through the post, of course, but there is no reason why outside doctors should not make more use, through this laboratory, of modern methods of diagnosis and treatment.

The following table shows the methods employed :—

| | 1923. | 1924. |
|--|-------|-------|
| Bacteriological and Protozoological— | | |
| Microscopical examinations | 577 | 705 |
| Agglutination tests | 147 | 237 |
| Preparation of vaccines | 67 | 75 |
| Decomplementising serum | 51 | 5 |
| Sigma re-action | 52 | 67 |
| Cultural examinations | 201 | 145 |
| Examinations of water supplies | 9 | 37 |
| Antiseptic co-efficients | 26 | ... |
| Biologic tests | 1 | 3 |
| Helminthological— | | |
| Microscopical examinations | 76 | 84 |
| Entomological— | | |
| Identification of insects | 1 | 2 |
| Miscellaneous— | | |
| Preparation of rat specimens | ... | 88 |
| Pathological— | | |
| Microscopical examinations | 113 | 151 |
| Sections of tumours, etc. | 25 | 61 |
| Chemical— | | |
| Tests | 33 | 26 |
| Quantitative estimations | 47 | 14 |
| Biologic tests | ... | ... |
| Medico-Legal— | | |
| Microscopical or chemical tests | 28 | 11 |
| Biologic tests | 2 | 3 |
| | <hr/> | <hr/> |
| | 1,456 | 1,714 |

II.—REMARKS ON DISEASES, ETC., DEALT WITH.

A. *Blackwater Fever*.—Unfortunately no research investigator could be sent out from England, and our own experiments on haemolysis could not be continued for the reasons already given, but a fresh start will now be made both by our new research investigator and by ourselves.

B. *Malaria*.—Three hundred and sixty-two examinations were made, with 38 positive results (33 malignant tertian distributed throughout the year, 2 benign tertian in June and August, and 3 quartan in January and December). It will be seen how often a single blood smear was negative, although many of the above cases were probably malarial (but with parasites too scanty to show up in a single smear). In negative cases a differential count is made, or further smears examined till a result is obtained, but it is obvious that some improvement on the single-smear method is desirable.

For purposes of research a series of malarial slides was examined, but the present differential count could not be improved upon. To report the presence of actual parasites is the ideal to be aimed at, and we intend to do further work this year on methods of concentrating the parasites, so that more positive results will be obtained.

C. *Undulant (Malta) Fever*.—One hundred and forty-two agglutination tests were made, with 26 positive results (of which several were only weakly positive and were reported as "doubtful"), as opposed to 19 positives last year. The disease seems fairly stationary.

As normal people and those suffering from other diseases often give a weak positive test, the question of diagnosis is of importance. Our method of blood culture is of help in this respect, and under research a new test was worked out which it is hoped will enable us to say definitely whether a patient has the disease or not. Some cultures of our strains were handed to Mr. Bevan for experimental purposes, but he was, unfortunately, unable to carry out the projected inoculations.

D. *Infection with Alcaligenes Fæcalis*.—As this organism is related to *Alcaligenes melitensis* and *A. abortus* of Malta fever, for purposes of research a series of 98 (suspected Malta fever) cases were specially tested against this organism as well, to see whether infection with it occurs in Rhodesia. Of the 98 cases, 5 gave positive results, 6 gave results too doubtful to call positive, and the rest were negative. Comparing the results with those obtained in the same cases against *melitensis* and *abortus*, we get the following table:—

| Group. | Results with | | | Number of cases. |
|-----------|--------------|----------|----------|------------------|
| | Melitensis. | Abortus. | Fæcalis. | |
| 1 | + | - | - | 2 |
| 2 | + | + | - | 9 |
| 3 | + | + | + | 2 |
| 4 | + | - | + | 1 |
| 5 | - | - | + | 2 |
| 6 | - | - | - | 82 |
| Total ... | | | | 98 |

Groups 1 and 2 were undoubtedly Malta fever. Groups 3 and 4 gave much stronger results in titre for both *melitensis* and *abortus* than for *fæcalis*, and were therefore accepted as cases of Malta fever, but the results show how closely the three organisms are connected. Group 5 can be accepted as cases of infection with *fæcalis*, especially as they were also negative for typhoid and para-typhoid as well. Group 6 were negative for all three organisms. Some of them were cases of enteric.

E. *Enteric (Typhoid) and Para-Typhoid Fevers*.—Eighty-eight agglutination tests gave 17 positive results for typhoid and 1 positive

for para-typhoid. Some blood cultures were also made. Other organisms causing similar disease are kept in stock, and are tested against when necessary.

Blood and stool cultures are not asked for as frequently as is advisable, as they often give a result in the first week of the diseases (while the agglutination test is still negative), and thus give earlier and consequently more valuable information.

F. *Fevers of Doubtful Causation.*—It will be seen that, out of a large number of cases examined for malaria, Malta and enteric fevers, only a few gave positive results. Most of the negative cases were probably malarial, as a negative for enteric or Malta fever can usually be accepted as such, while a negative for malaria is not so reliable; but there remain a certain number of cases in which the diagnosis is doubtful, and these are of the following types:—

(1) Cases giving a negative agglutination test for typhoid, para-typhoid and Malta fevers. This generally means of course that the patient has not got these diseases, but some other infection which has next to be ascertained; but in a few cases of typhoid and Malta the agglutination test is negative, either because the infection is so slight that no re-action (giving agglutination) takes place, or the infection is so severe that the patient is overwhelmed and no re-action can be produced. In such cases we have to repeat the test or do cultures, but this means delay in diagnosis. It is clear that, by means of research, we should aim at some test or improvements that will give an earlier and correct diagnosis, and this point is being worked upon.

(2) Cases which are doubtfully (because weakly) positive to the test, and yet are not suffering from the disease at all. Normal people, cases of pyrexia, previous inoculation, previous disease or similar disease may all give a positive test, and so cause doubt in diagnosis. Further tests usually clear the matter up, but this again means delay. Research is again necessary for improvement of diagnostic methods, and it is hoped that our new test, described in the next paragraph, will prove of value in this direction.

(3) Cases which give a positive agglutination, say, for both typhoid and para-typhoid, or both typhoid and Malta fevers. As double infections are rare, these cases are usually ones of single infection which happen to agglutinate the other organism because it is similar, or some of the causes mentioned in the second paragraph may be at work. It is often difficult to say which is the infecting organism, and it may require several tests to decide this, which all means delay. Under research, therefore, we devised last year a new test, called the saturation test, which gives promising results and will be further tested and reported on this year. It is briefly as follows:—If an organism be saturated with a heated allied (but not its own) agglutinating serum, it will be found that it can now no longer be agglutinated by the allied serum, but can still be agglutinated by its own serum. Thus a case of, say, typhoid will by this test give a positive for typhoid only, and cannot give a positive for any other disease.

I have gone rather fully into the above points to make it clear how important the question of early and accurate diagnosis is in such common and important diseases as are mentioned above. There is a tendency to regard research as only called for in a few, perhaps rare or striking diseases, but it is obvious that it is equally important in such common

diseases as malaria, enteric and Malta fevers, for nothing is more important than that both doctor and patient should know the diagnosis as early as possible.

G. *Dysentery (Bacillary and Protozoal)*.—Only two examinations were made for bacilli, although these are a common cause; Morgan's bacillus (an occasional cause) was found in one. Forty-four examinations were made for amoebæ, with 9 positive results.

H. *Pneumonia and Pneumococcal Infections*.—Pneumonia is now, apparently, well under control, owing to preventive inoculation and other measures, and no serious outbreak required investigation. The pneumococcus is, however, a common invader of the lungs and throat in other conditions, for which reason we usually have to include it in our vaccines for respiratory diseases.

I. *Influenza*.—No large epidemic has called for study. The filtrable virus (*B. pneumosintes*) is being studied, chiefly in America. A similar filtrable virus, belonging to the same group, occurs in horse-sickness, and has been studied by Lister at Johannesburg. Similar organisms occur in the Pasteurelloses (haemorrhagic septicæmias) of animals, and it is probable that all the above diseases, with perhaps pneumonic plague, are closely connected. This would explain the severity and haemorrhagic character of pandemic influenza. It is interesting to note that pneumonic plague, like influenza, is infective through the sputum chiefly, unlike bubonic plague, which is conveyed by fleas.

J. *Plague*.—Three examinations were made for the presence of *B. pestis*, with negative results. Eighty-eight rats of different kinds were sent to us for preparation as specimens of the different plague-bearing rats.

K. *Tuberculosis*.—Eighty examinations, with 17 positives, as opposed to 74 examinations, with 16 positives, in 1923. Most of these were lung cases. The disease seems fairly stationary in this Colony, though increase of population (through importation of outside cases) may cause some increase.

L. *Leprosy*.—Four negative examinations, as opposed to 3 positives in 1923.

M. *Diphtheria*.—Ten examinations, with 2 positives, as opposed to 56 examinations, with 18 positives, in 1923. A few positives are really chance infections, and not actually cases of the disease. The Salisbury Municipality are to be congratulated on taking steps, through their medical officer of health, towards giving the Schick test a trial, as it should prove of value in stamping out epidemics or in preventing their occurrence.

N. *Syphilis*.—Seventy examinations, chiefly Sigma tests, were made, with 22 positives. Where the Wasserman re-action is required we undertake to decomplementise and post the serum to Johannesburg, but we ourselves prefer the Sigma test, as equally reliable, simpler and perhaps more delicate. It is gratifying to note that the local doctors apparently agree with this opinion, for in 1924 there were 5 Wassermans and 67 Sigmas asked for, as opposed to 51 Wassermans and 52 Sigmas in 1923. We have asked doctors to try the Luetin test as well, as this is of value where the usual tests are negative, and I am glad to hear that this is being done.

O. *Gonorrhœa*.—Fifty-one examinations, with 18 positives and 2 suspicious results. The symptoms are quite often due to other organisms, such as straphylococci, streptococci and even diphtheroid bacilli.

P. *Schistosomiasis (Bilharzia)*.—Sixty-three examinations, with 21 positives for *S. hæmatobium* and 1 positive for *S. mansoni*. The disease does not seem to have increased at all since last year. Under research some snails from the Mazoe Dam were examined; of 8 Limnææ 3 were found infected with liver fluke of cattle, and of 3 Planorbis 1 showed cercariæ of immature growth. Physopsis is the usual carrier of *S. hæmatobium* and Limnææ only occasionally so, while Planorbis carries *S. mansoni*. Research into methods of combating the disease, such as by destruction of snails, is necessary. The latest methods of treatment are giving good results. There seems to be little danger in swimming baths, if properly protected by settling and filtration.

Q. *Diabetes*.—Urine is tested in connection with the Insulin treatment, and other necessary tests such as the blood-sugar test are made when required.

R. *Trypanosomiasis*.—Thirty-four examinations were made of a few patients, and 2 were found infected—one a European at the Lonely Mine, the other a native in Salisbury.

S. *Water Supplies*.—Thirty-seven examinations, as opposed to 9 in 1923, were made of the local water supplies from dam, wells, boreholes, municipal park and tobacco experiment station. It will be seen that the Salisbury Municipality, by frequent tests, ensure that a pure supply is given to the public.

T. *Pathology*.—One hundred and fifty-one examinations of deposits, etc., and 61 sections of tumours, etc., were made; 17 malignant growths, 5 non-malignant, 20 inflammatory conditions, 3 cysts, 1 hydatid, and 1 syphilitic condition were found.

U. *Chemical*.—Twenty chemical tests and 14 quantitative estimations were made.

V. *Medico-Legal*.—Eleven microscopical or chemical tests and 3 biologic tests were made with blood stains, etc., in police cases.

III.—STATEMENT OF FEES EARNED.

The fees classed below as "Government" represent work done for patients in hospital, police, paupers and natives. "Private" represents work done for private patients outside hospital.

Pasteur Institute—

| | 1923. | 1924. |
|----------------------|---------------|---------------|
| Private | £32 8 0 | £13 8 0 |
| Government | ... | ... |
| Total | <hr/> £32 8 0 | <hr/> £13 8 0 |

Public Health Laboratory—

| | | |
|----------------------|----------------|-----------------|
| Private | £272 18 9 | £379 17 6 |
| Government | 421 8 6 | 469 13 0 |
| Total | <hr/> £694 7 3 | <hr/> £809 10 6 |

Combined Totals—

| | | |
|----------------------|-----------------|-----------------|
| Private | £305 6 9 | £393 5 6 |
| Government | 421 8 6 | 469 13 0 |
| Total | <hr/> £726 15 3 | <hr/> £862 18 6 |

L. J. JOHN ORPEN,

Government Bacteriologist.

**Annual Report of the Medical Superintendent,
Ingutsheni Mental Hospital.**

I have the honour to present my report for the year ended 31st December, 1924.

On 1st January there were 174 patients on the register. During the year 70 were admitted, 22 were discharged, 2 were transferred to the Union, and 16 died. Two hundred and forty-four cases were treated, that is, 40 male Europeans, 171 male natives and 33 female natives. There remained in residence on the 31st December, 1924, 204 patients, *i.e.*, 30 Europeans, 142 male natives and 32 female natives.

The 70 cases admitted included 4 re-admissions, 3 Europeans and 1 native. Of the discharges, excluding transfers, 21 were discharged recovered and 1 relieved. The recovery rate per centum, calculated on the number of admissions, was 31.42, as against 30.76 for the previous year. The death rate, calculated on the number treated, was 6.55, as against 12.94 in 1923.

Of the two cases transferred, one, a male native, was sent to Pretoria. He belonged to the Transvaal and had been detained in this hospital as a criminal lunatic since March, 1917. Having incidentally mentioned this alien, I may refer to the increasing number of alien natives admitted. They add considerably to the inconvenience of overcrowding here, as well as being costly to maintain. The other case transferred to the Union was a male European who was acutely hallucinated and had developed homicidal tendencies. As there are no means of treating recalcitrant cases separately, it became necessary to remove him to Valkenberg, where, I am glad to state, he made a good recovery.

Seclusion and restraint were employed in the case of four Europeans and one native for periods ranging from five to thirty-eight hours on seven occasions. The reason for restraint was for violence to patients and staff. Seclusion was used to avoid struggling with patients suffering from maniacal excitement.

The general health was satisfactory. In August and September there was an outbreak of influenza, but it was of a mild form and only one death (a male native) resulted. Chicken-pox was introduced by a native visitor, but it was confined to the native female division, where almost all the inmates were infected.

Additional accommodation has been long promised, and is, I understand, shortly to be commenced. All through the year we had to contend with unexampled overcrowding, and in order to obtain relief we had to convert a store into sleeping quarters for natives. Further accommodation is urgently required, and this can only be obtained by an early start with the building programme. A feature of overcrowding is that direct admissions have to be received into dormitories. Patients frequently come in who are excited and violent, and who, for other reasons also, should be placed in single rooms; but, owing to the deficiency of this kind of accommodation, they have to be placed in dormitories, where they disturb the other patients and often cause and receive injuries.

Instances of complaints from relatives of female European mental patients regarding delay in arranging transfer from their homes to a

hospital in the Union have been brought to my notice, and I sincerely hope that accommodation for this class will be provided before the end of the current year.

Farming and garden operations are being extended yearly, the main drawback to constant supplies being shortage of water during the dry season. In common with other parts, the maize crop was a failure, but cotton, which was tried as an experiment, did rather well, and besides proving useful and congenial employment for native patients, the yield was encouraging; consequently a larger area has been put under cotton this season. The value of last year's crop has not yet been ascertained, but it should be about £100 net.

In order to keep this institution in good working order it may be necessary to obtain one or two men with training in mental nursing. We have already benefited by bringing two senior men from England. The men available locally are usually well on towards middle life; they have been, in many instances, earning big wages, and it is found that they are inclined to grumble at the irksomeness of the work and other conditions inseparable from institution life.

Revenue from paying patients and sales amounted to £1,039 4s. 7d., *i.e.*, maintenance fees £941 1s., sales £98 3s. 7d. Supplies from the garden and farm for the use of the hospital amounted in value to £753 3s. 10d. There were outstanding from maintenance fees and sales £233 9s. 3d. on the 31st December, 1924.

The total expenditure for the year, including value of produce supplied from farm and garden, was £5,977 5s. 8d. This works out at £31 10s. per patient per annum, as against £40 4s. 7d. for the previous year. The cost of maintenance, calculated on the gross expenditure, is 1s. 8½d.; and the cost per patient per diem, excluding produce, is 1s. 6d. The net cost to the Government, after deducting revenue from hospital vote of expenditure, is 1s. 2½d. per caput per diem. The amount per head shows an economical working, and is in part, at all events, due to the increase in the number of patients, and that there were supplies of mealie meal from the previous year's crop to meet requirements for nine months.

I cannot conclude this report without placing on record the deep sense of the loss which we have sustained in the death of Dr. W. M. Eaton, who was for so many years the superintendent of this hospital. He took the deepest interest in all that concerned the welfare of both patients and staff, and in the general development of the institution which he had charge of since its foundation.

A. W. FORRESTER,
Medical Superintendent.

Habit-forming Drugs.

Two alterations in the "Southern Rhodesia Opium and Habit-forming Drugs Regulations Proclamation, 1923," were made during the year.

Mention was made in the report for 1923 of the difficulty encountered in administering the regulations in so far as they related to habit-forming drugs required for veterinary purposes. This difficulty has now been overcome by the introduction of the following clauses as additional paragraphs to section 5, sub-section (2) of the aforesaid Proclamation.

"1. Any farmer or owner of live stock desiring to purchase from time to time tincture of opium (laudanum), for the prevention or treatment of disease in stock, may apply in writing to the Chief Veterinary Surgeon or any Government veterinary surgeon for a permit therefor, stating the quantity required, the approximate number of the various classes of stock kept and the person from whom it is proposed to purchase the drug.

2. The Chief Veterinary Surgeon or any Government veterinary surgeon may at his discretion, and after any enquiries which he may deem necessary, issue a permit accordingly.

3. Every such permit shall be subject to the following:—

- (a) the drug may be purchased only from the person specified in the permit and in quantity not exceeding 20 ounces at any one time;
- (b) the permit must be produced on the occasion of each purchase; the person from whom the purchase is made shall enter thereon the quantity supplied and the date of supply;
- (c) the holder of the permit shall not have in his possession more than 20 ounces of tincture of opium (laudanum) at any one time;
- (d) the container, in addition to any other label required by law, shall be labelled 'for animal treatment only,' and shall be kept under lock and key by the owner or some responsible person acting on his behalf, and the contents shall be used for administration to animals;
- (e) the permit shall remain in force and effect up to the end of the calendar year in which it is issued, but not longer; every applicant for a renewal shall return, with his application, the permit previously issued;
- (f) the Chief Veterinary Surgeon may cancel or withdraw any such permit at any time, and on being notified of such cancellation or withdrawal the holder shall forthwith return the permit to the Chief Veterinary Surgeon;
- (g) the holder of the permit shall produce the same for inspection, together with any tincture of opium in his possession, on the request of any person authorised thereto in writing by the Chief Veterinary Surgeon, or to any police officer of or above the rank of sergeant.

4. Copies of all such permits shall be rendered to the Chief Veterinary Surgeon, and a return of the same shall be rendered annually to the Medical Director, showing the number of permits issued, with the names of the persons to whom such permits have been granted, the district in which each such person resides and the amount of tincture of opium (laudanum) allowed to be purchased in each case."

The second alteration referred to section 5 of the regulations published under Government Notice No. 422 of 21st September, 1923, whereunder the list of preparations enumerated in Schedule "D" were excluded from the Proclamation and regulations. The preparations referred to were not similarly excluded elsewhere, and to bring the regulations into line with the legal provision existing elsewhere the following words were added to section 5 above referred to by Government Notice No. 476 of the 29th of August, 1924 :—"except as regards the importation into or exportation from Southern Rhodesia of such preparations." The absence of this provision was an obvious defect in the regulation as it existed formerly, and the amendment puts the matter in order.

Import certificates issued during the year numbered 41 and export certificates numbered 8.

Permits were issued to import the following habit-forming drugs :—

| | | |
|--------------------------|--------|--------|
| Cocaine hydrochloride | 15,005 | grains |
| Morphine | 12,569 | grains |
| Morphine sulphate | 7,152 | grains |
| Heroin hydrochloride | 5,377 | grains |
| Codein phosphate | 960 | grains |
| Morphine hydrochloride | 5,134 | grains |
| Cocaine | 42 | grains |
| Cocaine salicylate | 30 | grains |
| Opium | 750 | grains |
| Extract of opium | 150 | grains |
| Powdered opium | 2 | lbs. |
| Gall. and opium ointment | 7 | lbs. |
| Extract of Indian hemp | 960 | grains |
| Hemp seed | 4 | cwt. |

The export permits issued covered the following :—

| | | |
|---------------------------|-------|--------|
| Liquor opii sedativa | 20 | ounces |
| Morphine tartrate | 240 | grains |
| Morphine | 75 | grains |
| Cocaine hydrochloride | 16 | grains |
| Morphine hydrochloride | 30 | grains |
| Cocaine | 10 | grains |
| Atropine | 2 | grains |
| Scopolamine hydrochloride | 1 1/5 | grains |

The permits issued by the Veterinary Department under the provisions of Government Notice No. 368 of 27th June, 1924, numbered 11, and the total quantity of tincture of opium authorised to be purchased under the permits so granted was 132 ounces.

Seizures during the Year.—About one-sixteenth of an ounce of opium and an opium smoking outfit were seized at the Chinese resting house at Bulawayo. It was impossible, however, to prove ownership; no criminal action resulted therefore, and the articles seized were destroyed.

PART III.

TABLE 1.

EUROPEAN BIRTHS, 1924.

| District. | Jan. | Feb. | Mar. | April | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | Total. |
|---------------|------|------|------|-------|-----|------|------|------|-------|------|------|------|--------|
| Salisbury ... | 21 | 23 | 20 | 22 | 21 | 19 | 15 | 28 | 18 | 21 | 22 | 28 | 258 |
| Bulawayo ... | 24 | 21 | 17 | 23 | 31 | 28 | 19 | 26 | 17 | 21 | 19 | 16 | 262 |
| Umtali ... | 8 | 8 | 9 | 9 | 1 | 3 | 10 | 6 | 6 | 3 | 10 | 6 | 79 |
| Gwelo ... | 3 | 1 | 2 | 4 | 4 | 5 | 5 | 5 | 5 | 6 | 4 | 9 | 53 |
| Fort Victoria | 1 | 8 | 6 | 4 | 3 | 2 | 2 | 3 | ... | 5 | 3 | ... | 37 |
| Gatooma ... | 3 | 6 | 4 | 6 | 5 | 6 | 5 | 4 | 1 | 4 | 7 | 4 | 55 |
| Gwanda ... | ... | ... | 2 | ... | 2 | ... | 2 | 1 | 1 | ... | ... | ... | 8 |
| Selukwe ... | 1 | ... | 1 | 1 | ... | ... | 1 | 3 | 1 | 3 | 1 | ... | 12 |
| Charter ... | ... | 2 | 3 | 2 | 2 | 3 | 4 | 3 | 5 | 2 | 4 | 4 | 34 |
| Melsetter ... | 1 | 2 | ... | 2 | 4 | 4 | ... | 1 | 1 | 3 | 2 | 2 | 22 |
| Umvuma ... | 1 | 2 | ... | 1 | 4 | ... | 4 | ... | 2 | 5 | 2 | ... | 21 |
| Hartley ... | ... | ... | ... | ... | 1 | 3 | 1 | 1 | 1 | ... | 1 | ... | 8 |
| Que Que ... | 4 | 2 | 2 | 3 | 3 | 5 | ... | ... | ... | 4 | 1 | 1 | 25 |
| Totals ... | 67 | 75 | 66 | 77 | 81 | 78 | 68 | 81 | 58 | 77 | 76 | 70 | 874 |

TABLE 2.

EUROPEAN DEATHS, 1924.

| Age periods. | | | | Males. | Females. | Totals. |
|----------------|-----|-----|-----|--------|----------|---------|
| 0—1 | ... | ... | ... | 39 | 26 | 65 |
| 1—5 | ... | ... | ... | 10 | 8 | 18 |
| 5—15 | ... | ... | ... | 11 | 6 | 17 |
| 15—25 | ... | ... | ... | 7 | 13 | 20 |
| 25—35 | ... | ... | ... | 19 | 14 | 33 |
| 35—45 | ... | ... | ... | 16 | 9 | 25 |
| 45—55 | ... | ... | ... | 34 | 18 | 52 |
| 55—65 | ... | ... | ... | 30 | 6 | 36 |
| 65—75 | ... | ... | ... | 12 | 14 | 26 |
| 75—85 and over | ... | ... | ... | 7 | 8 | 15 |
| Age unknown | ... | ... | ... | 3 | ... | 3 |
| All ages | ... | ... | ... | 188 | 122 | 310 |

TABLE 3.
EUROPEAN DEATHS, 1924.

| District. | Jan. | Feb. | Mar. | April | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | Totals |
|---------------|------|------|------|-------|-----|------|------|------|-------|------|------|------|--------|
| Salisbury ... | 9 | 18 | 13 | 10 | 6 | 9 | 8 | 9 | 5 | 10 | 11 | 13 | 121 |
| Bulawayo ... | 9 | ... | 10 | 6 | 9 | 6 | 11 | 8 | 7 | 14 | 4 | 5 | 89 |
| Umtali ... | 1 | ... | 4 | 2 | ... | 2 | ... | 4 | 1 | 1 | 3 | ... | 18 |
| Gwelo ... | 1 | 2 | ... | 1 | 1 | 2 | 2 | 4 | 3 | 3 | 2 | 2 | 23 |
| Fort Victoria | 1 | 2 | ... | 3 | 1 | 3 | 2 | 2 | 1 | ... | 1 | 1 | 17 |
| Gatooma ... | ... | 1 | ... | ... | 2 | 1 | ... | 1 | ... | 3 | ... | ... | 8 |
| Gwanda ... | ... | ... | 1 | ... | ... | ... | 1 | 1 | ... | ... | ... | ... | 3 |
| Selukwe ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | ... | 2 | ... | 4 |
| Charter ... | 1 | ... | ... | ... | 1 | ... | ... | 2 | 1 | 1 | 2 | ... | 8 |
| Melsetter ... | ... | ... | ... | ... | 2 | 1 | ... | ... | 2 | ... | ... | ... | 5 |
| Umvuma ... | 1 | ... | 1 | ... | ... | 1 | 2 | 1 | 1 | ... | 1 | ... | 8 |
| Hartley ... | ... | ... | ... | ... | 1 | ... | 1 | ... | ... | ... | ... | ... | 2 |
| Que Que ... | ... | ... | ... | ... | 1 | 1 | ... | 1 | ... | 1 | ... | ... | 4 |
| Totals ... | 23 | 23 | 29 | 23 | 23 | 27 | 26 | 34 | 22 | 33 | 26 | 21 | 310 |

TABLE 4.
EUROPEAN BIRTHS AND DEATHS, 1924.

| Month. | Births. | Deaths. | Ages of the dying. | | | | | | | | | | |
|--------------------|---------|---------|--------------------|------|------|-------|-------|-------|-------|-------|-------|----------------------|---------------------|
| | | | 0-1 | 1-5 | 5-15 | 15-25 | 25-35 | 35-45 | 45-55 | 55-65 | 65-75 | 75-85 and over | Age un- known |
| January ... | 67 | 23 | 4 | ... | 1 | 1 | 2 | 3 | 2 | 3 | 4 | 2 | 1 |
| February | 75 | 23 | 7 | 2 | ... | 1 | 2 | 3 | 3 | 3 | 1 | 1 | ... |
| March ... | 66 | 29 | 3 | 3 | 1 | 2 | 5 | 3 | 6 | 3 | 2 | 1 | ... |
| April ... | 77 | 23 | 3 | ... | 4 | 3 | 1 | 3 | 5 | 2 | 1 | ... | 1 |
| May ... | 81 | 23 | 3 | ... | ... | 1 | 3 | 2 | 7 | 2 | 4 | 1 | ... |
| June ... | 78 | 27 | 6 | 2 | 2 | 2 | 3 | 1 | 3 | 3 | 5 | ... | ... |
| July ... | 68 | 26 | 8 | ... | ... | 3 | 1 | 1 | 5 | 6 | ... | 2 | ... |
| August ... | 81 | 34 | 8 | 2 | 1 | 3 | 5 | 1 | 7 | 4 | 2 | 1 | ... |
| September | 58 | 22 | 7 | 1 | ... | 1 | 3 | 1 | 3 | 4 | 2 | ... | ... |
| October ... | 77 | 33 | 7 | 3 | 2 | ... | 5 | 4 | 4 | 3 | 1 | 3 | 1 |
| November | 76 | 26 | 5 | 2 | 3 | 1 | 1 | 2 | 5 | 3 | 2 | 2 | ... |
| December | 70 | 21 | 4 | 3 | 3 | 2 | 2 | 1 | 2 | ... | 2 | 2 | ... |
| Totals ... | 874* | 310 | 65 | 18 | 17 | 20 | 33 | 25 | 52 | 36 | 26 | 15 | 3 |
| Per cent. of total | ... | 20.97 | 5.81 | 5.48 | 6.45 | 10.65 | 8.06 | 16.77 | 11.61 | 8.39 | 4.84 | .97 | |

26.77 per cent. of total European deaths.

* Illegitimate births 2.40 per cent. of total European births.

TABLE 5.

TABLE SHOWING EUROPEAN ADMISSIONS TO
HOSPITALS DURING 1924.

| Name of hospital. | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | Totals |
|-------------------|------|------|------|------|-----|------|------|------|-------|------|------|------|--------|
| Salisbury ... | 93 | 89 | 95 | 104 | 90 | 86 | 67 | 93 | 93 | 106 | 85 | 92 | 1,093 |
| Bulawayo ... | 70 | 38 | 66 | 63 | 54 | 78 | 51 | 54 | 58 | 70 | 66 | 63 | 731 |
| Umtali ... | 23 | 23 | 28 | 33 | 28 | 16 | 24 | 19 | 22 | 26 | 31 | 35 | 308 |
| Gwelo ... | 14 | 13 | 8 | 16 | 10 | 14 | 14 | 14 | 12 | 17 | 18 | 17 | 167 |
| Fort Victoria | 9 | 11 | 8 | 16 | 9 | 14 | 3 | 2 | 9 | 9 | 9 | 5 | 104 |
| Gatooma ... | 14 | 15 | 32 | 18 | 25 | 11 | 18 | 13 | 23 | 22 | 14 | 17 | 222 |
| Enkeldoorn | 3 | 2 | 4 | ... | ... | 1 | ... | ... | ... | 2 | 1 | ... | 13 |
| Gwanda ... | 3 | 3 | 1 | 3 | 6 | 2 | 3 | 3 | 3 | 6 | 5 | 4 | 42 |
| Sinoia ... | 4 | 6 | 5 | 9 | 5 | 4 | 10 | 4 | 3 | 4 | 5 | 8 | 67 |
| Shamva ... | 6 | 8 | 12 | 14 | 8 | 3 | 7 | 4 | 4 | 4 | 9 | 1 | 80 |
| Belingwe ... | 1 | ... | ... | ... | 1 | ... | ... | 2 | ... | ... | 1 | 1 | 6 |
| Totals ... | 240 | 208 | 259 | 276 | 236 | 229 | 197 | 208 | 227 | 266 | 244 | 243 | 2,833 |

TABLE 6.

TABLE SHOWING NATIVE ADMISSIONS TO
HOSPITALS DURING 1924.

| Name of hospital. | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | Totals |
|-------------------|------|------|------|------|-----|------|------|------|-------|------|------|------|--------|
| Salisbury ... | 78 | 81 | 87 | 86 | 77 | 71 | 77 | 86 | 114 | 94 | 96 | 82 | 1,029 |
| Bulawayo ... | 72 | 51 | 51 | 68 | 84 | 54 | 70 | 61 | 98 | 148 | 149 | 97 | 1,003 |
| Umtali ... | 14 | 15 | 12 | 15 | 28 | 15 | 18 | 24 | 32 | 20 | 38 | 54 | 285 |
| Gwelo ... | 55 | 45 | 44 | 48 | 38 | 36 | 23 | 31 | 43 | 91 | 61 | 50 | 565 |
| Fort Victoria | 10 | 8 | 7 | 16 | 4 | 12 | 9 | 8 | 8 | 9 | 9 | 9 | 109 |
| Gatooma ... | 37 | 28 | 44 | 31 | 41 | 37 | 33 | 46 | 47 | 40 | 52 | 36 | 472 |
| Enkeldoorn | 5 | 2 | 2 | 1 | 4 | 3 | 3 | 4 | 4 | 1 | 1 | 9 | 39 |
| Gwanda ... | 8 | 9 | 11 | 7 | 13 | 11 | 10 | 9 | 13 | 25 | 17 | 23 | 156 |
| Sinoia ... | 4 | 14 | 8 | 7 | 8 | 13 | 11 | 7 | 6 | 11 | 5 | 9 | 103 |
| Shamva ... | 15 | 15 | 16 | 19 | 20 | 20 | 9 | 17 | 16 | 7 | 10 | 12 | 176 |
| Belingwe ... | 9 | 3 | 3 | 10 | 2 | 2 | 2 | 4 | 5 | 2 | 5 | 7 | 54 |
| Totals ... | 307 | 271 | 285 | 308 | 319 | 274 | 265 | 297 | 386 | 448 | 443 | 388 | 3,991 |

TABLE 7.

Table showing monthly admissions to hospitals during 1924 from malaria, blackwater fever, dysentery, pneumonia, typhoid fever and scurvy.

EUROPEANS.

| Disease | Jan. | Feb. | Mar. | April | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | Totals |
|------------------|------|------|------|-------|-----|------|------|------|-------|------|------|------|--------|
| Malaria | 35 | 44 | 62 | 78 | 49 | 20 | 18 | 20 | 13 | 23 | 20 | 31 | 413 |
| Blackwater fever | 2 | 1 | 2 | 3 | 8 | 1 | 2 | ... | ... | 1 | ... | ... | 20 |
| Dysentery | 2 | 2 | 3 | ... | 2 | 1 | 1 | 2 | 4 | 10 | 3 | 5 | 35 |
| Pneumonia | 4 | 2 | 1 | 6 | 8 | 5 | 9 | 10 | 11 | 3 | 1 | 2 | 62 |
| Typhoid | 13 | 7 | 3 | 1 | 1 | 1 | ... | ... | 6 | 6 | 3 | 1 | 42 |
| Scurvy | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

NATIVES.

| | | | | | | | | | | | | | |
|------------------|-----|-----|-----|-----|----|-----|-----|-----|-----|-----|-----|-----|-----|
| Malaria | 30 | 32 | 24 | 44 | 28 | 31 | 12 | 10 | 15 | 35 | 20 | 19 | 300 |
| Blackwater fever | ... | ... | 1 | ... | 1 | ... | ... | ... | 1 | ... | ... | ... | 3 |
| Dysentery | ... | 3 | 1 | 4 | 3 | 2 | 6 | 4 | 8 | 5 | 3 | 6 | 45 |
| Pneumonia | 21 | 20 | 14 | 21 | 21 | 31 | 21 | 33 | 58 | 54 | 55 | 32 | 381 |
| Typhoid | ... | ... | 2 | ... | 1 | ... | ... | ... | ... | 4 | ... | 1 | 8 |
| Scurvy | 11 | 6 | ... | 6 | 3 | 7 | ... | 6 | 12 | 18 | 28 | 19 | 116 |

TABLE 8.

Cases, with mortality rate per cent., admitted to hospitals during 1924, as compared with 1923.

| Name of hospital | | 1923 | | | 1924 | | | |
|------------------|-----|--------|--------|--------------------------|-------|--------|--------------------------|-------|
| | | Cases | Deaths | Mortality rate per cent. | Cases | Deaths | Mortality rate per cent. | |
| Salisbury | ... | White | 1,072 | 32 | 2.99 | 1,093 | 49 | 4.48 |
| | | Native | 808 | 70 | 8.66 | 1,029 | 139 | 13.51 |
| Umtali | ... | White | 463 | 14 | 3.02 | 308 | 9 | 2.92 |
| | | Native | 137 | 13 | 9.49 | 285 | 11 | 3.86 |
| Gwelo | ... | White | 218 | 5 | 2.29 | 167 | 10 | 6.00 |
| | | Native | 400 | 41 | 10.25 | 565 | 62 | 10.97 |
| Fort Victoria | ... | White | 98 | 4 | 4.08 | 104 | 5 | 4.81 |
| | | Native | 110 | 10 | 9.09 | 109 | 11 | 10.09 |
| Gwanda | ... | White | 36 | 1 | 2.78 | 42 | 1 | 2.38 |
| | | Native | 124 | 9 | 7.26 | 156 | 22 | 14.10 |
| Enkeldoorn | ... | White | 32 | 3 | 9.37 | 13 | ... | ... |
| | | Native | 24 | 1 | 4.17 | 39 | 1 | 2.56 |
| Gatooma | ... | White | 262 | 14 | 5.34 | 222 | 7 | 3.15 |
| | | Native | 414 | 75 | 18.12 | 472 | 79 | 16.74 |
| Bulawayo | ... | White | 925 | 36 | 3.89 | 731 | 24 | 3.28 |
| | | Native | 682 | 59 | 8.65 | 1,003 | 85 | 8.47 |
| Shamva | ... | White | 125 | 6 | 4.80 | 80 | 2 | 2.50 |
| | | Native | 202 | 26 | 12.87 | 176 | 34 | 19.32 |
| Sinoia | ... | White | 73 | 2 | 2.74 | 67 | 2 | 3.00 |
| | | Native | 82 | 13 | 15.85 | 103 | 15 | 14.56 |
| Belingwe | ... | White | 6 | ... | ... | 6 | ... | ... |
| | | Native | 32 | 2 | 6.25 | 54 | 3 | 5.56 |
| Totals | | White | 3,310 | 117 | 3.53 | 2,833 | 109 | 3.85 |
| | | Native | 3,015 | 319 | 10.58 | 3,991 | 462 | 11.58 |

TABLE 9.

Cases, with mortality rate per cent., of malarial fever admitted to hospitals in 1924, as compared with 1923.

| Name of hospital. | | 1923. | | | 1924. | | | |
|-------------------|--------|---------------|------------|--------------------------|-------------|------------|--------------------------|-------------|
| | | Cases. | Deaths. | Mortality rate per cent. | Cases. | Deaths. | Mortality rate per cent. | |
| Salisbury | White | 175 | — | — | 113 | 4 | 3.54 | |
| | Native | 129 | — | — | 104 | 2 | 1.92 | |
| Umtali | White | 292 | 1 | 0.34 | 110 | — | — | |
| | Native | 27 | 1 | 3.70 | 29 | 1 | 3.45 | |
| Gwelo | White | 71 | 2 | 2.82 | 17 | — | — | |
| | Native | 53 | 5 | 9.43 | 44 | — | — | |
| Fort Victoria | White | 21 | — | — | 26 | 1 | 3.85 | |
| | Native | 9 | — | — | 10 | — | — | |
| Gwanda | White | 10 | — | — | 4 | — | — | |
| | Native | 11 | — | — | 7 | — | — | |
| Enkeldoorn | White | 6 | 1 | 16.67 | — | — | — | |
| | Native | 2 | — | — | 1 | — | — | |
| Gatooma | White | 106 | 3 | 2.83 | 44 | 1 | 2.27 | |
| | Native | 30 | 5 | 16.67 | 11 | — | — | |
| Bulawayo | White | 170 | 6 | 3.53 | 54 | — | — | |
| | Native | 56 | 2 | 3.57 | 59 | 1 | 1.69 | |
| Shamva | White | 70 | — | — | 32 | 1 | 3.12 | |
| | Native | 28 | 1 | 3.57 | 17 | 1 | 5.88 | |
| Sinoia | White | 28 | 1 | 3.57 | 11 | — | — | |
| | Native | 6 | — | — | 11 | — | — | |
| Belingwe | White | 4 | — | — | 2 | — | — | |
| | Native | 8 | — | — | 7 | — | — | |
| Totals | | White | 953 | 14 | 1.47 | 413 | 7 | 1.69 |
| | | Native | 359 | 14 | 3.90 | 300 | 5 | 1.67 |

TABLE 10.

Cases, with mortality rate per cent., of haemoglobinuric fever (blackwater) admitted to hospitals in 1924, as compared with 1923.

| Name of hospital. | | 1923. | | | 1924. | | | |
|-------------------|--------|---------------|-----------|--------------------------|--------------|-----------|--------------------------|--------------|
| | | Cases. | Deaths. | Mortality rate per cent. | Cases. | Deaths. | Mortality rate per cent. | |
| Salisbury | White | 12 | 5 | 41.67 | 3 | — | — | |
| | Native | — | — | — | 2 | 1 | 50.00 | |
| Umtali | White | 15 | 5 | 33.33 | 4 | — | — | |
| | Native | — | — | — | — | — | — | |
| Gwelo | White | 1 | — | — | — | — | — | |
| | Native | — | — | — | — | — | — | |
| Fort Victoria | White | 4 | — | — | 2 | 1 | 50.00 | |
| | Native | — | — | — | — | — | — | |
| Gwanda | White | 1 | 1 | — | — | — | — | |
| | Native | — | — | — | — | — | — | |
| Enkeldoorn | White | 1 | 1 | 100.00 | — | — | — | |
| | Native | — | — | — | — | — | — | |
| Gatooma | White | 10 | — | — | 2 | — | — | |
| | Native | 1 | — | — | — | — | — | |
| Bulawayo | White | 6 | 1 | 16.67 | 5 | — | — | |
| | Native | — | — | — | — | — | — | |
| Shamva | White | 12 | 2 | 16.67 | 3 | — | — | |
| | Native | — | — | — | — | — | — | |
| Sinoia | White | 1 | — | — | 1 | — | — | |
| | Native | — | — | — | 1 | 1 | 100.00 | |
| Belingwe | White | — | — | — | — | — | — | |
| | Native | — | — | — | — | — | — | |
| Totals | | White | 63 | 14 | 22.22 | 20 | 1 | 5.00 |
| | | Native | 1 | — | — | 3 | 2 | 66.67 |

TABLE 11.

Cases, with mortality rate per cent., of pneumonia admitted to hospitals during 1924, as compared with 1923.

| Name of hospital. | | 1923. | | | 1924. | | |
|-------------------|---|------------|---------|--------------------------|--------|---------|--------------------------|
| | | Cases. | Deaths. | Mortality rate per cent. | Cases. | Deaths. | Mortality rate per cent. |
| Salisbury | - | White 5 | — | — | 16 | 4 | 25.00 |
| | | Native 82 | 22 | 26.83 | 141 | 45 | 28.37 |
| Umtali | - | White 7 | — | — | 11 | 3 | 27.27 |
| | | Native 6 | — | — | 7 | 1 | 14.29 |
| Gwelo | - | White 2 | — | — | 10 | 1 | 10.00 |
| | | Native 14 | 8 | 57.14 | 63 | 17 | 27.00 |
| Fort Victoria | - | White 2 | — | — | 1 | — | — |
| | | Native 8 | 5 | 52.50 | 4 | 2 | 50.00 |
| Gwanda | - | White — | — | — | — | — | — |
| | | Native 3 | 2 | 66.67 | 14 | 7 | 50.00 |
| Enkeldoorn | - | White 4 | 1 | 25.00 | — | — | — |
| | | Native — | — | — | 1 | — | — |
| Gatooma | - | White 3 | — | — | 13 | 1 | 7.54 |
| | | Native 42 | 13 | 30.95 | 74 | 24 | 32.43 |
| Bulawayo | - | White 17 | 2 | 11.76 | 6 | 3 | 50.00 |
| | | Native 24 | 10 | 41.67 | 39 | 10 | 25.64 |
| Shamva | - | White 2 | 1 | 50.00 | 2 | — | — |
| | | Native 9 | 7 | 77.78 | 18 | 8 | 44.44 |
| Sinoia | - | White 1 | 1 | 100.00 | 2 | — | — |
| | | Native 6 | 2 | 33.33 | 12 | 3 | 25.00 |
| Belingwe | - | White — | — | — | 1 | — | — |
| | | Native 2 | — | — | 8 | — | — |
| Totals | | White 43 | 5 | 11.63 | 62 | 12 | 19.35 |
| | | Native 196 | 69 | 35.20 | 381 | 117 | 30.71 |

TABLE 12.

Cases, with mortality rate per cent., of dysentery admitted to hospitals during 1924, as compared with 1923.

| Name of hospital. | | 1923. | | | 1924. | | |
|-------------------|---|-----------|---------|--------------------------|--------|---------|--------------------------|
| | | Cases. | Deaths. | Mortality rate per cent. | Cases. | Deaths. | Mortality rate per cent. |
| Salisbury | - | White 6 | — | — | 21 | — | — |
| | | Native 7 | 3 | 42.68 | 10 | 2 | 20.00 |
| Umtali | - | White 2 | 1 | 50.00 | 1 | — | — |
| | | Native 4 | 1 | 25.00 | 2 | — | — |
| Gwelo | - | White 1 | — | — | — | — | — |
| | | Native 7 | 2 | 28.57 | 4 | 1 | 25.00 |
| Fort Victoria | - | White 6 | — | — | 3 | — | — |
| | | Native 1 | — | — | — | — | — |
| Gwanda | - | White — | — | — | 3 | — | — |
| | | Native — | — | — | 4 | — | — |
| Enkeldoorn | - | White — | — | — | — | — | — |
| | | Native — | — | — | — | — | — |
| Gatooma | - | White 2 | — | — | 1 | — | — |
| | | Native 4 | 2 | 50.00 | 1 | — | — |
| Bulawayo | - | White 6 | 1 | 16.67 | 2 | — | — |
| | | Native 6 | 1 | 16.67 | — | — | — |
| Shamva | - | White 2 | — | — | 2 | — | — |
| | | Native 9 | 1 | 11.11 | 16 | 3 | 18.75 |
| Sinoia | - | White 4 | — | — | 2 | — | — |
| | | Native 2 | 1 | 50.00 | 4 | 1 | 25.00 |
| Belingwe | - | White — | — | — | — | — | — |
| | | Native 1 | — | — | 4 | — | — |
| Totals | | White 29 | 2 | 6.90 | 35 | — | — |
| | | Native 41 | 11 | 26.83 | 45 | 7 | 15.56 |

TABLE 13.

Cases, with mortality rate per cent., of typhoid fever admitted to hospitals in 1924, as compared with 1923.

| Name of hospital. | | 1923. | | | 1924. | | | |
|-------------------|---|--------|---------|--------------------------|--------|---------|--------------------------|-------|
| | | Cases. | Deaths. | Mortality rate per cent. | Cases. | Deaths. | Mortality rate per cent. | |
| Salisbury | - | White | 7 | 1 | 14.29 | 7 | 1 | 14.29 |
| | | Native | 3 | 1 | 33.33 | 3 | — | — |
| Umtali | - | White | 3 | — | — | 1 | — | — |
| | | Native | — | — | — | — | — | — |
| Gwelo | - | White | 5 | — | — | 9 | 1 | 11.11 |
| | | Native | — | — | — | 4 | 1 | 25.00 |
| Fort Victoria | - | White | 1 | 1 | 100.00 | 2 | — | — |
| | | Native | — | — | — | — | — | — |
| Gwanda | - | White | 1 | — | — | — | — | — |
| | | Native | — | — | — | — | — | — |
| Enkeldoorn | - | White | — | — | — | 1 | — | — |
| | | Native | — | — | — | — | — | — |
| Gatooma | - | White | 7 | 2 | 28.57 | 3 | 2 | 66.67 |
| | | Native | — | — | — | — | — | — |
| Bulawayo | - | White | 26 | 3 | 11.52 | 18 | 2 | 11.11 |
| | | Native | 7 | 3 | 42.86 | 1 | — | — |
| Shamva | - | White | 1 | — | — | 1 | — | — |
| | | Native | 1 | — | — | — | — | — |
| Sinoia | - | White | — | — | — | — | — | — |
| | | Native | — | — | — | — | — | — |
| Belingwe | - | White | — | — | — | — | — | — |
| | | Native | — | — | — | — | — | — |
| Totals | | White | 51 | 7 | 13.73 | 42 | 6 | 14.29 |
| | | Native | 10 | 4 | 40.00 | 8 | 1 | 12.50 |

TABLE 14.

Cases, with mortality rate per cent., of scurvy admitted to hospitals in 1924, as compared with 1923.

| Name of hospital. | | 1923. | | | 1924. | | | |
|-------------------|---|--------|---------|--------------------------|--------|---------|--------------------------|-------|
| | | Cases. | Deaths. | Mortality rate per cent. | Cases. | Deaths. | Mortality rate per cent. | |
| Salisbury | - | White | — | — | — | — | — | |
| | | Native | — | — | — | 14 | 1 | 7.14 |
| Umtali | - | White | — | — | — | — | — | — |
| | | Native | 1 | — | — | — | — | — |
| Gwelo | - | White | — | — | — | — | — | — |
| | | Native | 21 | 1 | 4.76 | 7 | 3 | 42.86 |
| Fort Victoria | - | White | — | — | — | — | — | — |
| | | Native | — | — | — | — | — | — |
| Gwanda | - | White | — | — | — | — | — | — |
| | | Native | 1 | — | — | 17 | 1 | 5.88 |
| Enkeldoorn | - | White | — | — | — | — | — | — |
| | | Native | — | — | — | — | — | — |
| Gatooma | - | White | — | — | — | — | — | — |
| | | Native | 28 | 3 | 10.71 | 20 | — | — |
| Bulawayo | - | White | — | — | — | — | — | — |
| | | Native | 8 | 2 | 25.00 | 57 | 5 | 8.70 |
| Shamva | - | White | — | — | — | — | — | — |
| | | Native | — | — | — | — | — | — |
| Sinoia | - | White | — | — | — | — | — | — |
| | | Native | 3 | — | — | — | — | — |
| Belingwe | - | White | — | — | — | — | — | — |
| | | Native | 1 | — | — | 1 | — | — |
| Totals | | White | — | — | — | — | — | — |
| | | Native | 63 | 6 | 9.52 | 116 | 10 | 8.62 |

TABLE I5.—RETURN OF DEATHS REGISTERED DURING THE YEAR 1924.

EUROPEANS.

| Name of disease. | EUROPEANS. | | | | | | | | | | | | NATIVES. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|------------|-----|-----|-----|-----|-----|-----|-----|------|-----|-----|-----|----------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|-----|-------|-----|----|---|-------|--|--|--|-------------|--|--|--|--------|--|--|--|
| | 0-1 | | | | 1-5 | | | | 5-15 | | | | 15-25 | | | | 25-35 | | | | 35-45 | | | | 45-55 | | | | 55-65 | | | | 65-75 | | | | 75-85 | | | | Age unknown | | | | Totals | | | |
| | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | | | | | | | | | | | | |
| I. General Diseases. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1. Typhoid fever | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 3 | | | | | | | | | | | | | |
| 3. Relapsing fever | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 7 | | | | | | | | | | | | | |
| 4. Malaria | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 4 | | | | | | | | | | | | | |
| 4a. Blackwater fever | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 8 | | | | | | | | | | | | | |
| 8. Whooping cough | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 3 | | | | | | | | | | | | | |
| 10. Influenza | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 8 | | | | | | | | | | | | | |
| 14. Dysentery | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 10 | | | | | | | | | | | | | |
| 20. Purulent infection and septicæmia | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 4 | | | | | | | | | | | | | |
| 24. Tetanus | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 3 | | | | | | | | | | | | | |
| 26. Pellagra | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 27. Beri-beri | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 4 | | | | | | | | | | | | | |
| 28. Tuberculosis of the lungs | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 81 | | | | | | | | | | | | | |
| 29. Acute miliary tuberculosis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 30. Tuberculous meningitis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 31. Abdominal tuberculosis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 2 | | | | | | | | | | | | | |
| 32. Potts' disease | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 34. Tuberculosis of other organs | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 35. Disseminated tuberculosis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 37. Syphilis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 11 | | | | | | | | | | | | | |
| 39. Cancer and other malignant tumours of the buccal cavity | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 40. Cancer and other malignant tumours of the stomach, liver | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 3 | | | | | | | | | | | | | |
| 41. Cancer and other malignant tumours of the peritoneum, intestines, rectum | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 42. Cancer and other malignant tumours of the female genital organs | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 2 | | | | | | | | | | | | | |
| 43. Cancer and other malignant tumours of the breast | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 44. Cancer and other malignant tumours of the skin | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 45. Cancer and other malignant tumours of other organs, or of organs not specified | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 46. Other tumours (tumours of the female genital organs excepted) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 47. Acute articular rheumatism | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 48. Chronic rheumatism and gout | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 49. Scurvy | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 9 | | | | | | | | | | | | | |
| 50. Diabetes | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 53. Leucæmia | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| 54. Anæmia, chlorosis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | | | | | | | | | | | | | |
| Carried forward | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 49 | | | | | | | | | | | | | |
| Grand totals | 6 | 5 | 5 | 4 | 4 | 3 | 4 | 5 | 11 | 4 | 8 | 3 | 10 | 7 | 10 | 2 | 7 | 6 | 2 | 2 | ... | ... | 67 | 41 | 2 | 3 | 162 | 5 | 164 | 8 | 231 | 49 | | | | | | | | | | | | | | | | |

EUROPEANS.

| Name of disease. | EUROPEANS. | | | | | | | | | | | | NATIVES. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|------------|-----|-----|-----|-----|-----|-----|-----|------|-----|-----|-----|----------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|-----|-------|-----|-----|--|-------------|--|--|--|--------|--|--|--|
| | 0-1 | | | | 1-5 | | | | 5-15 | | | | 15-25 | | | | 25-35 | | | | 35-45 | | | | 45-55 | | | | 55-65 | | | | 65-75 | | | | 75-85 | | | | Age unknown | | | | Totals | | | |
| | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | | | | | | | | | | | | |
| Brought forward | ... | ... | 20 | 16 | 9 | 7 | 6 | 3 | 7 | 10 | 14 | 6 | 14 | 4 | 23 | 14 | 23 | 5 | 10 | 11 | 4 | 4 | 1 | ... | 131 | 80 | 10 | 14 | 357 | 17 | 367 | 31 | 498 | 111 | | | | | | | | | | | | | | |
| 105. Diarrhoea and enteritis (two years and over) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | |
| 108. Appendicitis and typhilitis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | |
| 109. Hernia, intestinal obstruction | ... | ... | 1 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 113. Cirrhosis of the liver | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 115. Other diseases of the liver | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 116. Diseases of the spleen | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 117. Simple peritonitis (non-puerperal) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| VI. Non-Venereal Diseases of the Genito-Urinary System and Annexa. | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 119. Acute nephritis | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 122. Other diseases of the kidneys and annexa | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 124. Diseases of the bladder | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 127. Non-venereal diseases of the male genital organs | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 130. Other diseases of the uterus | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 131. Cysts and other tumours of the ovary | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 132. Salpingitis and other diseases of the female genital organs | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| VII. The Puerperal State. | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 134. Accidents of pregnancy | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 135. Puerperal hemorrhage | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 136. Other accidents of labour | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 137. Puerperal septicæmia | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 138. Puerperal albuminuria and convulsions | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| VIII. Diseases of the Skin and of the Cellular Tissue. | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 142. Gangrene | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 144. Acute abscess | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 145. Other diseases of the skin and anæxia | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| IX. Diseases of the Bones and of the Organs of Locomotion. | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 146. Diseases of the bones (tuberculosis excepted) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| 147. Diseases of the joints (tuberculosis and rheumatism excepted) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | | | | | | | | | | |
| Carried forward | ... | ... | 22 | 16 | 9 | 7 | 9 | 4 | 7 | 13 | 15 | 13 | 14 | 9 | 26 | 18 | 25 | 5 | 10 | 11 | 4 | 5 | 1 | ... | 142 | 101 | 10 | 14 | 392 | 23 | 402 | 37 | 544 | 138 | | | | | | | | | | | | | | |

EUROPEANS.

| Name of disease. | EUROPEANS. | | | | | | | | | | | | NATIVES. | | | | | | | | | | | | Grand totals | | | | | | | | | |
|--|------------|-----|-----|-----|-----|-----|------|-----|-----|-------|-----|-----|----------|-----|-----|-------|-----|-----|-------|-----|----|-------|-----|-----|--------------|-----|-----|-------|-----|-----|-----|-----|-----|-----|
| | 0—1 | | | 1—5 | | | 5—15 | | | 15—25 | | | 25—35 | | | 35—45 | | | 45—55 | | | 55—65 | | | 65—75 | | | 75—85 | | | | | | |
| | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | M | F | | | | |
| Brought forward | ... | ... | 22 | 16 | 9 | 7 | 9 | 4 | 7 | 13 | 15 | 13 | 14 | 9 | 26 | 18 | 25 | 5 | 10 | 11 | 4 | 5 | 1 | ... | 142 | 101 | 10 | 14 | 392 | 23 | 402 | 37 | 544 | 138 |
| X. Malformations. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 150. Congenital malformations (still-births not included) | ... | 1 | ... | ... | 1 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 2 | ... | ... | ... | ... | ... | ... | ... | ... | ... | 2 | ... | | |
| XI. Early Infancy. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 151. Congenital debility, icterus and sclerema | ... | ... | 16 | 10 | 1 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 17 | 10 | 4 | 1 | ... | ... | 4 | 1 | 21 | 11 | 3 | | | |
| XII. Old Age. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 154. Senility | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 2 | 1 | 3 | 2 | ... | ... | 5 | 3 | ... | 7 | ... | 7 | ... | 12 | 3 | |
| XIII. External Causes. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 155. Suicide by poison | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 157. Suicide by hanging and strangulation | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 159. Suicide by firearms | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 160. Suicide by cutting and piercing instruments | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 163. Other suicides | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 166. Conflagration | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 168. Absorption of deleterious gases (conflagration excepted) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 170. Traumatism by firearms | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 173. Traumatism in mines and quarries | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 175. Traumatism by other crushing (vehicles, railways, landslides, etc.) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 185. Fractures (cause not specified) | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 186. Other external violence | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 186a. Execution | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| XIV. Ill-defined Diseases. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 187. Ill-defined organic disease | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 188. Sudden death | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| 189. Cause of death not specified or ill-defined | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | |
| Totals | ... | ... | 39 | 26 | 10 | 8 | 11 | 6 | 7 | 13 | 19 | 14 | 16 | 9 | 34 | 18 | 30 | 6 | 12 | 14 | 7 | 8 | 3 | ... | 188 | 122 | 17 | 16 | 436 | 25 | 453 | 41 | 641 | 163 |

TABLE 16.

CLASSIFICATION OF DEATHS (EUROPEANS), 1924.
Deaths classified according to the international classification of causes of sickness and death.

| Classification No. | Disease. | No. of Deaths. |
|--------------------|--|----------------|
| 1 | Typhoid fever ... | 8 |
| 3 | Relapsing fever ... | 1 |
| 4 | Malaria ... | 13 |
| 4a | Blackwater fever ... | 11 |
| 8 | Whooping cough ... | 8 |
| 10 | Influenza ... | 11 |
| 14 | Dysentery ... | 6 |
| 20 | Purulent infection and septicæmia ... | 7 |
| 28 | Tuberculosis of the lungs ... | 14 |
| 29 | Acute miliary tuberculosis ... | 1 |
| 35 | Disseminated tuberculosis ... | 1 |
| 39 | Cancer and other malignant tumours of the buccal cavity ... | 1 |
| 40 | Cancer and other malignant tumours of the stomach, liver ... | 4 |
| 41 | Cancer and other malignant tumours of the peritoneum, intestines, rectum ... | 4 |
| 42 | Cancer and other malignant tumours of the female genital organs ... | 1 |
| 43 | Cancer and other malignant tumours of the breast ... | 3 |
| 44 | Cancer and other malignant tumours of the skin ... | 1 |
| 45 | Cancer and other malignant tumours of other organs, or of organs not specified ... | 7 |
| 46 | Other tumours (tumours of the female genital organs excepted) ... | 1 |
| 47 | Acute articular rheumatism ... | 1 |
| 48 | Chronic rheumatism and gout ... | 1 |
| 50 | Diabetes ... | 1 |
| 53 | Leucæmia ... | 1 |
| 54 | Anæmia, chlorosis ... | 1 |
| 55 | Other general diseases ... | 3 |
| 56 | Alcoholism ... | 3 |
| 60 | Encephalitis ... | 2 |
| 61 | Simple meningitis ... | 3 |
| 61c | Cerebro-spinal fever ... | 1 |
| 64 | Cerebral haemorrhage—apoplexy ... | 14 |
| 66 | Paralysis without specified cause ... | 1 |
| 68 | Other forms of mental alienation ... | 2 |
| 69 | Epilepsy ... | 2 |
| 71 | Convulsions of infants ... | 2 |
| 74 | Other diseases of the nervous system ... | 1 |
| 76 | Diseases of the ears ... | 3 |
| 77 | Pericarditis ... | 1 |
| 78 | Acute endocarditis ... | 2 |
| 79 | Organic diseases of the heart ... | 14 |
| 80 | Angina pectoris ... | 4 |
| 85 | Haemorrhage; other diseases of the circulatory system ... | 1 |
| 86 | Diseases of the nasal fossæ ... | 1 |
| 89 | Acute bronchitis ... | 4 |
| 90 | Chronic bronchitis ... | 1 |
| 91 | Broncho-pneumonia ... | 6 |
| 92 | Pneumonia ... | 20 |
| 93 | Pleurisy ... | 1 |
| 96 | Asthma ... | 2 |
| 98a | Miners' phthisis ... | 1 |
| 101 | Diseases of the oesophagus ... | 1 |
| 104 | Diarrhoea and enteritis (under two years) ... | 7 |
| 105 | Diarrhoea and enteritis (two years and over) ... | 1 |
| 108 | Appendicitis and typhilitis ... | 4 |
| 109 | Hernia, intestinal obstruction ... | 2 |
| 113 | Cirrhosis of the liver ... | 1 |
| 115 | Other diseases of the liver ... | 1 |
| 116 | Diseases of the spleen ... | 1 |
| 117 | Simple peritonitis (non-puerperal) ... | 4 |
| 119 | Acute nephritis ... | 4 |
| 122 | Other diseases of the kidneys and annexa ... | 1 |
| 130 | Other diseases of the uterus ... | 1 |
| 132 | Salpingitis and other diseases of the female genital organs ... | 1 |
| 134 | Accidents of pregnancy ... | 1 |
| 135 | Puerperal haemorrhage ... | 1 |
| 136 | Other accidents of labour ... | 1 |
| 137 | Puerperal septicæmia ... | 5 |
| 142 | Gangrene ... | 2 |
| 146 | Diseases of the bones (tuberculosis excepted) ... | 1 |
| 150 | Congenital malformations (still births not included) ... | 2 |
| 151 | Congenital debility, icterus and sclerema ... | 27 |
| 154 | Senility ... | 8 |
| 155 | Suicide by poison ... | 5 |
| 159 | Suicide by firearms ... | 4 |
| 163 | Other suicides ... | 1 |
| 166 | Conflagration ... | 1 |
| 173 | Traumatism in mines and quarries ... | 2 |
| 175 | Traumatism by other crushing (vehicles, railways, landslides, etc.) ... | 2 |
| 185 | Fractures (cause not specified) ... | 3 |
| 186 | Other external violence ... | 1 |
| 187 | Ill-defined organic disease ... | 1 |
| 188 | Sudden death ... | 2 |
| 189 | Cause of death not specified or ill-defined ... | 8 |
| Total | ... | 310 |

TABLE 17.

CLASSIFICATION OF DEATHS (NATIVES AND COLOURED), 1924.

Deaths classified according to the international classification of causes of sickness and death.

| Classification No. | Disease. | No. of Deaths. |
|--------------------|--|----------------|
| 1 | Typhoid fever ... | 2 |
| 4 | Malaria ... | 8 |
| 4a | Blackwater fever ... | 1 |
| 8 | Whooping cough ... | 1 |
| 10 | Influenza ... | 35 |
| 14 | Dysentery ... | 7 |
| 20 | Purulent infection and septicæmia ... | 10 |
| 24 | Tetanus ... | 3 |
| 26 | Pellagra ... | 1 |
| 27 | Beri-beri ... | 1 |
| 28 | Tuberculosis of the lungs ... | 71 |
| 30 | Tuberculous meningitis ... | 1 |
| 31 | Abdominal tuberculosis ... | 2 |
| 32 | Potts' disease ... | 2 |
| 34 | Tuberculosis of other organs ... | 1 |
| 35 | Disseminated tuberculosis ... | 1 |
| 37 | Syphilis ... | 11 |
| 45 | Cancer and other malignant tumours of other organs, or of organs not specified ... | 4 |
| 47 | Acute articular rheumatism ... | 1 |
| 49 | Scurvy ... | 9 |
| 55 | Other general diseases ... | 2 |
| 55a | Trypanosomiasis ... | 1 |
| 60 | Encephalitis ... | 3 |
| 61 | Simple meningitis ... | 22 |
| 61c | Cerebro-spinal fever ... | 1 |
| 63 | Other diseases of the spinal cord ... | 3 |
| 64 | Cerebral haemorrhage—apoplexy ... | 5 |
| 66 | Paralysis without specified cause ... | 4 |
| 68 | Other forms of mental alienation ... | 11 |
| 69 | Epilepsy ... | 6 |
| 76 | Diseases of the ears ... | 3 |
| 77 | Pericarditis ... | 1 |
| 78 | Acute endocarditis ... | 6 |
| 79 | Organic diseases of the heart ... | 6 |
| 81 | Diseases of the arteries, atheroma, aneurism, etc. ... | 1 |
| 82 | Embolism and thrombosis ... | 1 |
| 85 | Hæmorrhage; other diseases of the circulatory system ... | 2 |
| 89 | Acute bronchitis ... | 4 |
| 90 | Chronic bronchitis ... | 2 |
| 91 | Broncho-pneumonia ... | 7 |
| 92 | Pneumonia ... | 130 |
| 96 | Asthma ... | 2 |
| 104 | Diarrhoea and enteritis (under two years) ... | 3 |
| 105 | Diarrhoea and enteritis (two years and over) ... | 3 |
| 109 | Hernia, intestinal obstruction ... | 3 |
| 113 | Cirrhosis of the liver ... | 5 |
| 115 | Other diseases of the liver ... | 1 |
| 116 | Diseases of the spleen ... | 5 |
| 117 | Simple peritonitis (non-puerperal) ... | 3 |
| 119 | Acute nephritis ... | 2 |
| 124 | Diseases of the bladder ... | 1 |
| 127 | Non-venereal diseases of the male genital organs ... | 1 |
| 131 | Cysts and other tumours of the ovary ... | 1 |
| 132 | Salpingitis and other diseases of the female genital organs ... | 1 |
| 136 | Other accidents of labour ... | 1 |
| 137 | Puerperal septicæmia ... | 1 |
| 138 | Puerperal albuminuria and convulsions ... | 1 |
| 142 | Gangrene ... | 4 |
| 144 | Acute abscess ... | 4 |
| 145 | Other diseases of the skin and annexa ... | 2 |
| 146 | Diseases of the bones (tuberculosis excepted) ... | 1 |
| 147 | Diseases of the joints (tuberculosis and rheumatism excepted) ... | 1 |
| 151 | Congenital debility, icterus and sclerema ... | 5 |
| 154 | Senility ... | 7 |
| 157 | Suicide by hanging and strangulation ... | 1 |
| 159 | Suicide by firearms ... | 1 |
| 160 | Suicide by cutting and piercing instruments ... | 1 |
| 166 | Conflagration ... | 6 |
| 168 | Absorption of deleterious gases (conflagration excepted) ... | 1 |
| 170 | Traumatism by firearms ... | 2 |
| 173 | Traumatism in mines and quarries ... | 1 |
| 175 | Traumatism by other crushing (vehicles, railways, landslides, etc.) ... | 2 |
| 185 | Fractures (cause not specified) ... | 10 |
| 186 | Other external violence ... | 1 |
| 186a | Execution ... | 8 |
| 187 | Ill-defined organic disease ... | 1 |
| 189 | Cause of death not specified or ill-defined ... | 8 |
| Total | | 494 |

Included in the foregoing are the following Indian and coloured deaths, classified as follows:—

| Classification No. | Disease. | No. of Deaths. | |
|---|----------|----------------|-----------|
| | | Indian. | Coloured. |
| 4 Malaria | ... | 2 | 1 |
| 4a Blackwater fever | ... | — | 1 |
| 8 Whooping cough | ... | — | 1 |
| 10 Influenza | ... | 3 | — |
| 28 Tuberculosis of the lungs | ... | — | 3 |
| 37 Syphilis | ... | 1 | — |
| 47 Acute articular rheumatism | ... | 1 | — |
| 61 Simple meningitis | ... | — | 3 |
| 63 Other diseases of the spinal cord | ... | 1 | — |
| 69 Epilepsy | ... | — | 1 |
| 78 Acute endocarditis | ... | — | 1 |
| 79 Organic diseases of the heart | ... | 2 | 1 |
| 89 Acute bronchitis | ... | — | 1 |
| 91 Broncho-pneumonia | ... | — | 1 |
| 92 Pneumonia | ... | — | 1 |
| 96 Asthma | ... | 1 | — |
| 104 Diarrhoea and enteritis (under two years) | ... | 1 | 1 |
| 151 Congenital debility, icterus and sclerema | ... | 1 | — |
| 175 Traumatism by crushing (vehicles, railways, landslides, etc.) | ... | — | 1 |
| 189 Cause of death not specified or ill-defined | ... | 1 | 1 |
| Totals | ... | 14 | 18 |

TABLE 18.

Return of diseases and deaths (in-patients) in all Government hospitals and Memorial Hospital, Bulawayo, for the year 1924.

EUROPEANS.

| Diseases. | Remaining in hospital at end of 1923. | Yearly total. | | Total cases treated. | Remaining in hospital at end of 1924. |
|--------------------------------------|--|------------------|---------|----------------------------|--|
| | | Admis- sions. | Deaths. | | |
| INFECTIVE DISEASES. | | | | | |
| Dysentery—Amœbic | ... | 30 | ... | 30 | 1 |
| Bacillary | ... | 5 | ... | 5 | ... |
| Enteric | 7 | 42 | 6 | 49 | 2 |
| Erysipelas | ... | 5 | ... | 5 | ... |
| Gonorrhœa | ... | 1 | ... | 1 | ... |
| Influenza | ... | 91 | ... | 91 | 2 |
| Malaria—(a) Tertian | 3 | 161 | 5 | 164 | 4 |
| (c) Æstivo-autumnal | 1 | 244 | 2 | 245 | 3 |
| (d) Chronic | 1 | 8 | ... | 9 | 1 |
| (e) Blackwater | ... | 20 | 1 | 20 | 1 |
| Measles | ... | 1 | ... | 1 | ... |
| Malta fever | 3 | 16 | ... | 19 | ... |
| Pneumonia | 2 | 69 | 16 | 71 | ... |
| Relapsing fever | ... | 3 | ... | 3 | ... |
| Rheumatic fever | 1 | 9 | 1 | 10 | 1 |
| Septicæmia | ... | 6 | 6 | 6 | ... |
| Trypanosomiasis (sleeping sickness) | ... | 1 | ... | 1 | ... |
| Syphilis—(b) Secondary | ... | 2 | ... | 2 | ... |
| (c) Inherited | 1 | ... | ... | 1 | ... |
| Tetanus | ... | 1 | ... | 1 | ... |
| Tuberculosis | 5 | 20 | 9 | 25 | 2 |
| Other infective diseases | ... | 5 | ... | 5 | 2 |
| INTOXICATIONS. | | | | | |
| Alcoholism | 1 | 12 | ... | 13 | 1 |
| Others | ... | 1 | ... | 1 | ... |
| GENERAL DISEASES. | | | | | |
| Anæmia | ... | 12 | ... | 12 | ... |
| Anæmia—pernicious | ... | 1 | 1 | 1 | ... |
| Diabetes | ... | 1 | 1 | 1 | ... |
| Exophthalmic goitre | 1 | ... | ... | 1 | ... |
| Gout | 1 | 1 | ... | 2 | ... |
| Leucocytæmia | ... | 1 | 1 | 1 | ... |
| Other general diseases | 1 | 51 | ... | 52 | 2 |
| LOCAL DISEASES. | | | | | |
| Diseases of the nervous system— | | | | | |
| Sub-section 1— | | | | | |
| Neuritis | 4 | 28 | ... | 32 | ... |
| Meningitis | ... | 2 | 1 | 2 | ... |
| Myelitis | ... | 1 | ... | 1 | ... |
| Encephalitis | ... | 1 | 1 | 1 | ... |
| Abscess of brain | ... | 1 | 1 | 1 | ... |
| Congestion of brain | ... | 2 | ... | 2 | ... |
| Sub-section 2— | | | | | |
| Apoplexy | ... | 11 | 4 | 11 | 2 |
| Paralysis | 2 | 4 | ... | 6 | 1 |
| Chorea | ... | 1 | ... | 1 | ... |
| Epilepsy | 1 | 5 | ... | 6 | 1 |
| Neuralgia | ... | 4 | ... | 4 | ... |
| Hysteria | ... | 8 | ... | 8 | ... |
| Sub-section 3—mental diseases— | | | | | |
| Idiocy | ... | 2 | ... | 2 | ... |
| Dementia | 1 | ... | ... | 1 | ... |
| Other diseases of the nervous system | 3 | 35 | 2 | 38 | 2 |
| Diseases of the eye— | | | | | |
| Conjunctivitis | 2 | 13 | ... | 15 | ... |
| Keratitis | 1 | 3 | ... | 4 | ... |
| Ulceration of cornea | ... | 8 | ... | 8 | ... |
| Iritis | ... | 3 | ... | 3 | ... |
| Cataract | 1 | 1 | ... | 2 | 1 |
| Other diseases of the eye | ... | 15 | ... | 15 | ... |
| Carried forward | 43 | 968 | 58 | 1,011 | 29 |

| Diseases. | Remaining in hospital at end of 1923. | Yearly total. | | Total cases treated. | Remaining in hospital at end of 1924. |
|--|--|------------------|---------|----------------------------|--|
| | | Admis- sions. | Deaths. | | |
| Brought forward | 43 | 968 | 58 | 1,011 | 29 |
| Diseases of the ear— | | | | | |
| Inflammation | | 6 | ... | 6 | ... |
| Other diseases | | 4 | ... | 4 | ... |
| Diseases of the nose | | 10 | ... | 10 | ... |
| Diseases of the circulatory system— | | | | | |
| Pericarditis | | 4 | 2 | 4 | 1 |
| Endocarditis | | 8 | 2 | 8 | 1 |
| Valvular mitral | | 5 | ... | 5 | ... |
| Arterial sclerosis | | 7 | 1 | 7 | ... |
| Other diseases of the circulatory system ... | | 29 | 1 | 29 | 2 |
| Diseases of the respiratory system— | | | | | |
| Laryngitis | | 10 | ... | 10 | 1 |
| Bronchitis | | 47 | ... | 47 | ... |
| Broncho-pneumonia | | 12 | 3 | 12 | ... |
| Pleurisy | | 8 | ... | 8 | 2 |
| Empyema | | 3 | ... | 3 | ... |
| Other diseases of the respiratory system | | 15 | ... | 15 | ... |
| Diseases of the digestive system— | | | | | |
| Stomatitis | | 7 | ... | 7 | ... |
| Caries of teeth | | 22 | ... | 22 | ... |
| Glossitis | | 5 | ... | 5 | ... |
| Sore throat | | 7 | ... | 7 | ... |
| Inflammation of tonsils | | 3 | 70 | 73 | ... |
| Gastritis | | 1 | 29 | 30 | ... |
| Ulceration of stomach | | 1 | 9 | 9 | 1 |
| Hæmatemesis | | 1 | 1 | 2 | ... |
| Stricture of stomach | | 4 | 3 | 4 | ... |
| Dyspepsia | | 3 | ... | 3 | ... |
| Enteritis | | 18 | ... | 18 | 1 |
| Appendicitis | | 8 | 183 | 191 | 10 |
| Coitis | | 18 | 1 | 18 | ... |
| Ulceration of intestines | | 2 | ... | 2 | 1 |
| Hernia | | 1 | 29 | 30 | 1 |
| Diarrhoea | | 15 | ... | 15 | 1 |
| Constipation | | 1 | 9 | 10 | ... |
| Colic | | 17 | ... | 17 | ... |
| Haemorrhoids | | 1 | 17 | 18 | 1 |
| Hepatitis—acute | | 5 | ... | 5 | ... |
| Abscess of the liver | | 4 | ... | 4 | 1 |
| Cirrhosis | | 1 | 6 | 7 | 1 |
| Jaundice | | 1 | ... | 1 | ... |
| Peritonitis | | 2 | 6 | 8 | ... |
| Ascites | | ... | 2 | 2 | ... |
| Gallstones | | 18 | 2 | 18 | 2 |
| Other diseases of the digestive system ... | | 30 | ... | 30 | 1 |
| Diseases of the lymphatic system— | | | | | |
| Inflammation of lymphatic gland | | 1 | ... | 1 | ... |
| Diseases of the urinary system— | | | | | |
| Acute nephritis | | 6 | 1 | 6 | ... |
| Bright's disease | | 6 | 2 | 6 | ... |
| Pyelitis | | 5 | ... | 5 | ... |
| Calculus | | 4 | ... | 4 | ... |
| Renal colic | | 1 | 6 | 7 | ... |
| Cystitis | | 16 | 1 | 16 | 1 |
| Vesical calculus | | 2 | ... | 2 | ... |
| Hæmaturia | | 1 | ... | 1 | ... |
| Other diseases of the urinary system | | 4 | 1 | 4 | ... |
| Diseases of the generative system— | | | | | |
| Male organs— | | | | | |
| Urethritis | | 1 | ... | 1 | ... |
| Stricture | | 9 | ... | 9 | ... |
| Prostatitis | | 5 | ... | 5 | 1 |
| Hydrocele | | 3 | ... | 3 | ... |
| Orchitis | | 6 | ... | 6 | 1 |
| Epididymitis | | 2 | ... | 2 | 1 |
| Other diseases of the male organs ... | | 13 | ... | 13 | ... |
| Female organs— | | | | | |
| Ovaritis | | 6 | ... | 6 | ... |
| Ovarian cyst | | 8 | ... | 8 | ... |
| Endometritis | | 1 | 45 | 46 | ... |
| Displacement of uterus | | 2 | 8 | 10 | 1 |
| Vaginitis | | 1 | ... | 1 | ... |
| Dysmenorrhœa | | 5 | ... | 5 | ... |
| Menorrhagia | | 2 | ... | 2 | ... |
| Leucorrhœa | | 2 | ... | 2 | ... |
| Carried forward | 67 | 1,839 | 88 | 1,906 | 62 |

| Diseases. | Remaining in hospital at end of 1923. | Yearly total. | | Total cases treated. | Remaining in hospital at end of 1924. |
|---|--|------------------|---------|----------------------------|--|
| | | Admis- sions. | Deaths. | | |
| Brought forward | 67 | 1,839 | 88 | 1,906 | 62 |
| Diseases of the generative system (continued)— | | | | | |
| Female organs (continued)— | | | | | |
| Abortion | 1 | 53 | ... : : | 54 | ... |
| Retained placenta | ... | 7 | ... : : | 7 | ... |
| Premature birth | ... | 5 | ... : : | 5 | ... |
| Puerperal septicæmia | ... | 2 | 2 | 2 | ... |
| Mastitis | ... | 5 | 1 | 5 | ... |
| Abscess of breast | 1 | 2 | ... : : | 3 | ... |
| Other diseases of the female organs | 1 | 26 | ... : : | 27 | 2 |
| Confinements | 1 | 37 | ... : : | 38 | ... |
| Diseases of organs of locomotion— | | | | | |
| Osteitis | ... | 7 | 1 | 7 | ... |
| Arthritis | ... | 12 | ... : : | 12 | ... |
| Bursitis | ... | 3 | ... : : | 3 | ... |
| Other diseases of organs of locomotion ... | 1 | 14 | ... : : | 15 | ... |
| Diseases of connective tissue— | | | | | |
| Cellulitis | 2 | 41 | ... : : | 43 | 2 |
| Abscess | ... | 41 | ... : : | 41 | 2 |
| Elephantiasis | 1 | ... | ... : : | 1 | ... |
| Diseases of the skin— | | | | | |
| Urticaria | ... | 4 | ... : : | 4 | ... |
| Eczema | ... | 3 | ... : : | 3 | ... |
| Boil | ... | 9 | ... : : | 9 | ... |
| Carbuncle | ... | 11 | ... : : | 11 | 1 |
| Herpes | ... | 6 | ... : : | 6 | ... |
| Tinea | ... | 3 | ... : : | 3 | ... |
| Ulcers | ... | 17 | ... : : | 17 | 1 |
| Other diseases of the skin | ... | 12 | ... : : | 12 | ... |
| Injuries—general | | | | | |
| local | 5 | 132 | 3 | 137 | 6 |
| fractures | 12 | 112 | ... : : | 124 | ... |
| Surgical operations—major | ... | 53 | 1 | 53 | 4 |
| minor | ... | ... | ... : : | (471) | ... |
| Tumours | ... | ... | ... : : | (700) | ... |
| Poisons | 12 | 54 | 11 | 66 | 2 |
| Snake bite | ... | 10 | 2 | 10 | 1 |
| Parasites— | | | | | |
| Cestoda— <i>Tænia solium</i> | ... | 4 | ... : : | 4 | ... |
| <i>Tænia saginata</i> | ... | 1 | ... : : | 1 | ... |
| Not otherwise classified | ... | 244 | ... : : | 244 | ... |
| Gun-shot wounds | 1 | 12 | ... : : | 13 | ... |
| Bilharzia | ... | 10 | ... : : | 10 | ... |
| Senility | ... | 4 | ... : : | 4 | 3 |
| Not diagnosed | ... | 35 | ... : : | 35 | 5 |
| Burns | ... | 1 | ... : : | 1 | ... |
| Totals | 105 | 2,833 | 109 | 2,938 | 91 |

TABLE 19.

Return of diseases and deaths (in-patients) in all Government hospitals and Memorial Hospital, Bulawayo, for the year 1924.

NATIVES.

| Diseases. | Remaining in hospital at end of 1923. | Yearly total. | | Total cases treated. | Remaining in hospital at end of 1924. |
|--------------------------------------|--|------------------|---------|----------------------------|--|
| | | Admis- sions. | Deaths. | | |
| INFECTIVE DISEASES. | | | | | |
| Beri-beri | ... | 1 | 1 | 1 | ... |
| Diphtheria | ... | 4 | ... | 4 | ... |
| Dysentery—Amœbic | ... | 44 | 7 | 44 | 3 |
| Bacillary | ... | 1 | ... | 1 | ... |
| Enteric | ... | 8 | 1 | 8 | ... |
| Gonorrhœa | ... | 11 | ... | 11 | ... |
| Influenza | 8 | 332 | 30 | 340 | 8 |
| Leprosy—(a) Nodular | ... | 2 | ... | 2 | ... |
| Malaria—(a) Tertian | ... | 169 | 2 | 169 | ... |
| (b) Quartan | ... | 17 | 1 | 17 | ... |
| (c) Æstivo-autumnal | 6 | 114 | 2 | 120 | 6 |
| (d) Chronic | 1 | ... | ... | 1 | ... |
| (e) Blackwater | ... | 3 | 2 | 3 | ... |
| Pneumonia | 6 | 381 | 117 | 387 | 16 |
| Rheumatic fever | 2 | 30 | 1 | 32 | 1 |
| Septicæmia | ... | 10 | 8 | 10 | 1 |
| Trypanosomiasis (sleeping sickness) | 1 | 1 | 1 | 2 | 1 |
| Syphilis—(a) Primary | 5 | 90 | 3 | 95 | 12 |
| (b) Secondary | 5 | 40 | 5 | 45 | 1 |
| (c) Inherited | 1 | ... | ... | 1 | ... |
| Tetanus | ... | 3 | 3 | 3 | ... |
| Tuberculosis | 16 | 132 | 88 | 148 | 16 |
| Other infective diseases | ... | 2 | 1 | 2 | ... |
| GENERAL DISEASES. | | | | | |
| Anæmia | 1 | 8 | ... | 9 | 3 |
| Exophthalmic goitre | ... | 2 | ... | 2 | ... |
| Purpura | 2 | 10 | 1 | 12 | 1 |
| Scurvy | 7 | 116 | 10 | 123 | 26 |
| Other general diseases | 2 | 69 | 3 | 71 | 3 |
| LOCAL DISEASES. | | | | | |
| Diseases of the nervous system— | | | | | |
| Sub-section 1— | | | | | |
| Neuritis | 5 | 34 | ... | 39 | 3 |
| Meningitis | 1 | 31 | 26 | 32 | ... |
| Myelitis | ... | 1 | 1 | 1 | ... |
| Encephalitis | ... | 2 | 2 | 2 | ... |
| Abscess of brain | ... | 1 | ... | 1 | 1 |
| Sub-section 2— | | | | | |
| Apoplexy | ... | 11 | 5 | 11 | ... |
| Paralysis | 1 | 21 | 7 | 22 | 7 |
| Epilepsy | 2 | 17 | 3 | 19 | ... |
| Neuralgia | ... | 2 | ... | 2 | ... |
| Hysteria | ... | 2 | ... | 2 | ... |
| Sub-section 3—Mental Diseases— | | | | | |
| Idiocy | ... | 4 | ... | 4 | 1 |
| Mania | ... | 1 | ... | 1 | ... |
| Dementia | ... | 8 | 1 | 8 | ... |
| Other diseases of the nervous system | ... | 16 | ... | 16 | ... |
| Diseases of the eye— | | | | | |
| Conjunctivitis | ... | 67 | ... | 67 | 5 |
| Keratitis | ... | 7 | ... | 7 | ... |
| Ulceration of cornea | 2 | 12 | ... | 14 | ... |
| Iritis | ... | 6 | ... | 6 | ... |
| Optic neuritis | ... | 2 | ... | 2 | ... |
| Cataract | ... | 5 | ... | 5 | ... |
| Other diseases of the eye | ... | 8 | ... | 8 | ... |
| Diseases of the ear— | | | | | |
| Inflammation | 2 | 6 | ... | 8 | ... |
| Other diseases | ... | 5 | ... | 5 | ... |
| Diseases of the nose | 1 | ... | ... | 1 | ... |
| Carried forward | 77 | 1,869 | 332 | 1,926 | 115 |

| Diseases. | Remaining in hospital at end of 1923. | Yearly total. | | Total cases treated. | Remaining in hospital at end of 1924. |
|---|--|------------------|---------|----------------------------|--|
| | | Admis- sions. | Deaths. | | |
| Brought forward | 77 | 1,869 | 332 | 1,926 | 115 |
| Diseases of the circulatory system— | | | | | |
| Pericarditis | ... | 3 | 3 | 3 | ... |
| Endocarditis | 1 | 6 | 4 | 7 | ... |
| Valvular mitral | ... | 6 | 1 | 6 | ... |
| Arterial sclerosis | ... | 3 | 1 | 3 | 1 |
| Aneurism | ... | 1 | ... | 1 | ... |
| Other diseases of the circulatory system | 1 | 19 | 5 | 20 | 2 |
| Diseases of the respiratory system— | | | | | |
| Laryngitis | ... | 4 | ... | 4 | ... |
| Bronchitis | 4 | 48 | 7 | 52 | 2 |
| Broncho-pneumonia | ... | 25 | 4 | 25 | ... |
| Abscess of lung | ... | 1 | 1 | 1 | ... |
| Pleurisy | ... | 24 | ... | 24 | 1 |
| Other diseases of the respiratory system | ... | 7 | 2 | 7 | ... |
| Diseases of the digestive system— | | | | | |
| Stomatitis | 1 | 4 | ... | 5 | ... |
| Caries of teeth | ... | 1 | ... | 1 | ... |
| Glossitis | ... | 2 | ... | 2 | ... |
| Sore throat | 1 | 1 | ... | 2 | ... |
| Inflammation of tonsils | 1 | 25 | ... | 26 | ... |
| Gastritis | ... | 21 | ... | 21 | ... |
| Dyspepsia | ... | 2 | ... | 2 | ... |
| Enteritis | ... | 4 | 1 | 4 | ... |
| Appendicitis | 1 | 6 | ... | 7 | ... |
| Colitis | 1 | 2 | ... | 3 | ... |
| Ulceration of intestines | ... | 1 | 1 | 1 | ... |
| Hernia | 1 | 12 | 1 | 13 | ... |
| Diarrhoea | 1 | 14 | ... | 15 | ... |
| Constipation | 1 | 18 | ... | 19 | ... |
| Colic | ... | 6 | ... | 6 | ... |
| Hæmorrhoids | ... | 2 | ... | 2 | 1 |
| Pancreatitis | ... | 1 | 1 | 1 | ... |
| Hepatitis—acute | 1 | 1 | ... | 2 | ... |
| Abscess of the liver | ... | 2 | 1 | 2 | ... |
| Cirrhosis | ... | 7 | 5 | 7 | ... |
| Jaundice | ... | 2 | 2 | 2 | ... |
| Peritonitis | ... | 11 | 8 | 11 | 1 |
| Ascites | 3 | 9 | 2 | 12 | 1 |
| Other diseases of the digestive system ... | ... | 4 | 1 | 4 | 2 |
| Diseases of the lymphatic system— | | | | | |
| Splenitis | 1 | 7 | ... | 8 | 1 |
| Inflammation of lymphatic gland | ... | 8 | ... | 8 | 1 |
| Suppuration of lymphatic gland | ... | 1 | ... | 1 | ... |
| Lymphangitis | 3 | 61 | 1 | 64 | 6 |
| Other diseases of the lymphatic system ... | ... | 4 | 4 | 4 | ... |
| Diseases of the urinary system— | | | | | |
| Acute nephritis | ... | 4 | 1 | 4 | ... |
| Bright's disease | ... | 3 | 2 | 3 | ... |
| Pyelitis | ... | 2 | 1 | 2 | ... |
| Renal colic | ... | 2 | ... | 2 | ... |
| Cystitis | 1 | 3 | ... | 4 | ... |
| Vesical calculus | ... | 1 | ... | 1 | ... |
| Hæmaturia | ... | 3 | ... | 3 | ... |
| Other diseases of the urinary system ... | ... | 2 | ... | 2 | ... |
| Diseases of the generative system— | | | | | |
| Male organs— | | | | | |
| Urethritis | 3 | 1 | ... | 4 | ... |
| Gleet | ... | 1 | ... | 1 | ... |
| Stricture | ... | 3 | ... | 3 | 1 |
| Inflammation of scrotum | ... | 3 | ... | 3 | 1 |
| Hydrocele | ... | 6 | 1 | 6 | ... |
| Orchitis | 1 | 14 | ... | 15 | 2 |
| Other diseases of the male organs ... | ... | 3 | ... | 3 | ... |
| Female organs— | | | | | |
| Ovarian cyst | ... | 1 | 1 | 1 | ... |
| Endometritis | ... | 2 | ... | 2 | ... |
| Vaginitis | ... | 3 | ... | 3 | ... |
| Dysmenorrhœa | ... | 1 | ... | 1 | ... |
| Menorrhagia | ... | 2 | ... | 2 | ... |
| Leucorrhœa | ... | 1 | ... | 1 | ... |
| Abortion | ... | 4 | ... | 4 | ... |
| Delayed labour | ... | 3 | ... | 3 | ... |
| Retained placenta | ... | 4 | ... | 4 | ... |
| Premature birth | ... | 1 | ... | 1 | ... |
| Carried forward | 104 | 2,328 | 394 | 2,412 | 138 |

| Diseases. | Remaining in hospital at end of 1923. | Yearly total. | | Total cases treated. | Remaining in hospital at end of 1924. |
|---|--|------------------|---------|----------------------------|--|
| | | Admis- sions. | Deaths. | | |
| Brought forward | 104 | 2,328 | 394 | 2,412 | 138 |
| Diseases of the generative system (continued)— | | | | | |
| Female organs (continued)— | | | | | |
| Puerperal septicæmia | ... | 1 | ... | 1 | ... |
| Mastitis | ... | 2 | ... | 2 | ... |
| Abscess of breast | ... | 3 | ... | 3 | ... |
| Other diseases of the female organs | ... | 6 | 1 | 6 | 1 |
| Confinements | ... | 3 | 1 | 3 | ... |
| Diseases of organs of locomotion— | | | | | |
| Osteitis | 2 | 8 | 1 | 10 | |
| Arthritis | 1 | 21 | 2 | 22 | 2 |
| Spondylitis | 1 | 3 | ... | 4 | ... |
| Bursitis | ... | 10 | ... | 10 | ... |
| Other diseases of the organs of locomotion | ... | 14 | ... | 14 | ... |
| Diseases of connective tissue— | | | | | |
| Cellulitis | 2 | 105 | 2 | 107 | 9 |
| Abscess | 5 | 66 | 3 | 71 | 8 |
| Diseases of the skin— | | | | | |
| Urticaria | ... | 1 | ... | 1 | ... |
| Eczema | ... | 9 | ... | 9 | ... |
| Boil | ... | 6 | ... | 6 | ... |
| Psoriasis | ... | 1 | ... | 1 | ... |
| Oriental sore | 3 | 34 | ... | 37 | 10 |
| Tinea | ... | 3 | ... | 3 | ... |
| Scabies | ... | 5 | ... | 5 | ... |
| Acne | ... | 1 | ... | 1 | ... |
| Ulcers | 40 | 215 | 2 | 255 | 17 |
| Other diseases of the skin | 1 | 8 | ... | 9 | 3 |
| Injuries—general | | | | | |
| local | 23 | 503 | 21 | 526 | 42 |
| fractures | 64 | 287 | ... | 351 | 35 |
| Surgical operations—major | | | | | |
| minor | ... | ... | ... | (105) | ... |
| Tumours | | | | | |
| Cancer | ... | ... | ... | 74 | 6 |
| Malformations | 1 | ... | ... | (318) | ... |
| Poisons | ... | 2 | ... | 23 | ... |
| Snake bite | ... | 8 | ... | 1 | ... |
| Parasites— | | | | | |
| Cestoda— <i>Tænia solium</i> | 1 | 2 | ... | 3 | ... |
| <i>Tænia saginata</i> | ... | 2 | ... | 2 | ... |
| Not otherwise classified | ... | 114 | 11 | 114 | 1 |
| Gun-shot wound | 1 | 12 | 3 | 13 | 1 |
| Hydatids | ... | 1 | ... | 1 | ... |
| Bilharzia | ... | 10 | ... | 10 | ... |
| Senility | ... | 3 | 2 | 3 | 1 |
| Not diagnosed | ... | 19 | 1 | 19 | ... |
| Ill-defined | ... | 23 | 2 | 23 | ... |
| Burns | ... | 37 | 4 | 37 | 7 |
| Onyalai | ... | 2 | ... | 2 | ... |
| Suicide—cut throat | ... | 1 | 1 | 1 | ... |
| Totals | | | | | |
| | 250 | 3,991 | 462 | 4,241 | 283 |

TABLE 20.

Table giving the number of beds in each Government Hospital and Ingutsheni Mental Hospital, the daily average number of patients treated, the revenue and expenditure of each and the approximate charge on public funds for each patient in hospital during 1924.

| Name of hospital. | No. of beds. | | Daily average of patients treated. | | | No. of nursing staff. | No. of native staff. | Gross expenditure. | Revenue. | Deficit of revenue over expenditure. | £ s. d. | Approximate charge on public funds for each patient treated. |
|----------------------------|--------------|----------------------|------------------------------------|----------------------|-----------------------------------|-----------------------|----------------------|--------------------|----------|--------------------------------------|---------|--|
| | White. | Coloured and native. | White. | Coloured and native. | Total white, coloured and native. | | | | | | | |
| Salisbury | ... 60 | 100 | 47.9 | 64.4 | 112.3 | 36 | 53 | 17,748 | 6,839 | 10,909 | 4 18 6 | |
| Umtali ... | ... 36 | 20 | 11.5 | 16 | 27.5 | 6 | 13 | 3,900 | 1,158 | 2,742 | 4 10 2 | |
| Gwelo ... | ... 28 | 34 | 13.25 | 44.25 | 57.5 | 6 | 17 | 4,692 | 1,593 | 3,099 | 3 19 5 | |
| Fort Victoria | ... 13 | 13 | .41 | .63 | 1.04 | 3 | 10 | 1,706 | 329 | 1,377 | 6 3 6 | |
| Enkeldoorn | ... 4 | 4 | .53 | 2.22 | 2.75 | 1 | 3 | 705 | 27 | 678 | 12 6 7 | |
| Gwanda | ... 10 | 18 | 1.49 | 7.13 | 8.62 | 2 | 5 | 961 | 258 | 703 | 3 9 8 | |
| Gatooma | ... 16 | 60 | 5.9 | 70.4 | 76.3 | 6 | 15 | 4,116 | 1,448 | 2,668 | 3 9 8 | |
| Shamva | ... 14 | 15 | 1.56 | 11.32 | 12.88 | 3 | 12 | 1,944 | 364 | 1,580 | 5 19 1 | |
| Sinoia ... | ... 11 | 8 | 1.6 | 6.21 | 7.81 | 2 | 6 | 1,109 | 212 | 897 | 5 3 8 | |
| Belingwe ... | ... 9 | 15 | .14 | 5.11 | 5.25 | 1 | 2 | 509 | 59 | 450 | 7 10 0 | |
| Ingutsheni Mental Hospital | Mental 36 | 144 | 28.9 | 162.5 | 191.4 | 9 | 17 | 6,420 | 1,039 | 5,381 | 18 19 3 | |

TABLE 21.

STATEMENT OF PROGRESS AT GOVERNMENT HOSPITALS FOR THE YEAR 1924.

| Name of hospital. | Total number of patients maintained. | Total expenditure. | NO. OF UNITS MAINTAINED. | | | | | | | | EXPENDITURE | | | | | | | | | | | | | | | | EARNINGS. | | | | REVENUE RECEIVED. | | | | TOTAL AMOUNT OUTSTANDING. | | | | | |
|-------------------|--------------------------------------|--------------------|--------------------------|---------|-----------|---------|--------|---------|---|------|-------------|------|---|------|-------|-----|---|-----|-----|-----|------------------------------|------|--------|------|-----------|-----|-----------|-------|------------------|--------|-------------------|-------|---|------|---------------------------|----------|---------------------------|-----|-------|-------|
| | | | Staff. | | Patients. | | Total. | | Provisions and medical comforts, excluding produce. | | | | Drugs, surgical instruments and sundries. | | | | Furniture, equipment, clothing and repairs. | | | | Laundry staff and materials. | | | | Sanitary. | | | | Paying patients. | | | | Revenue per cent. of total expenditure. | | | | At end of preceding year. | | | |
| | | | White. | Native. | White. | Native. | White. | Native. | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | £ | | |
| Salisbury | ... 2,215 | 17,748 | 13,432 | 20,083 | 17,511 | 23,527 | 30,943 | 43,610 | 2,850 | 16.0 | 2,110 | 11.9 | 903 | 5.1 | 1,703 | 9.6 | 504 | 2.8 | 24 | .2 | 3,013 | 17.0 | 5,684 | 32.0 | 222 | 1.3 | 7,327 | 1,105 | 3,195 | 11,627 | 5,960 | 879 | 6,839 | 38.5 | 4 9 | 4 18 6 | 735 | 4.1 | 3,644 | 3,793 |
| Umtali | ... 608 | 3,900 | 2,337 | 4,285 | 4,209 | 5,834 | 6,546 | 10,119 | 453 | 11.6 | 482 | 12.4 | 510 | 13.0 | 160 | 4.1 | 163 | 4.2 | 101 | 2.6 | 612 | 15.7 | 1,263 | 32.4 | 30 | .8 | 1,352 | 227 | 813 | 2,392 | 1,013 | 145 | 1,158 | 29.7 | 4 8 | 4 10 2 | 126 | 3.2 | 222 | 493 |
| Gwelo | ... 781 | 4,692 | 2,558 | 5,124 | 4,837 | 16,121 | 7,395 | 21,245 | 624 | 13.3 | 533 | 11.4 | 341 | 7.3 | 246 | 5.2 | 125 | 2.7 | 192 | 4.1 | 950 | 20.2 | 1,541 | 32.8 | 14 | .3 | 898 | 938 | 1,792 | 3,628 | 691 | 902 | 1,593 | 33.9 | 3 4 | 3 19 5 | 126 | 2.7 | 319 | 554 |
| Victoria | ... 223 | 1,706 | 1,098 | 3,660 | 1,409 | 3,147 | 2,507 | 6,807 | 268 | 15.7 | 176 | 10.3 | 153 | 9.0 | 85 | 5.0 | 29 | 1.7 | 25 | 1.5 | 182 | 10.7 | 717 | 42.0 | 8 | .4 | 468 | 102 | 502 | 1,072 | 306 | 23 | 329 | 19.3 | 3 8 | 6 3 6 | 63 | 3.7 | 122 | 168 |
| Gwanda | ... 202 | 961 | 1,095 | 1,825 | 543 | 2,654 | 1,638 | 4,479 | 173 | 18.0 | 90 | 9.4 | 28 | 3.0 | 36 | 3.7 | 32 | 3.3 | ... | ... | 155 | 16.1 | 398 | 41.4 | 7 | .7 | 209 | 196 | 166 | 571 | 69 | 189 | 258 | 26.8 | 3 1 | 3 9 8 | 42 | 4.4 | 54 | 170 |
| Enkeldoorn | ... 55 | 705 | 732 | 1,098 | 195 | 813 | 927 | 1,911 | 81 | 11.5 | 52 | 7.4 | 53 | 7.5 | 30 | 4.3 | 7 | 1.0 | 18 | 2.6 | 89 | 12.6 | 353 | 50.0 | 1 | .1 | 10 | ... | 143 | 153 | 22 | 5 | 27 | 3.8 | 5 0 | 12 6 7 | 21 | 3.0 | 72 | 28 |
| Gatooma | ... 766 | 4,116 | 1,830 | 5,398 | 2,172 | 25,694 | 4,002 | 31,092 | 515 | 12.5 | 490 | 11.9 | 255 | 6.2 | 235 | 5.7 | 111 | 2.7 | 68 | 1.7 | 739 | 17.9 | 1,550 | 37.6 | 48 | 1.2 | 1,160 | 697 | 2,589 | 4,446 | 871 | 577 | 1,448 | 35.2 | 2 4 | 3 9 8 | 105 | 2.6 | 1,012 | 1,075 |
| Shamva | ... 267 | 1,944 | 1,166 | 4,392 | 569 | 4,133 | 1,735 | 8,525 | 288 | 14.8 | 217 | 11.2 | 134 | 6.9 | 67 | 3.4 | 53 | 2.7 | ... | ... | 249 | 12.8 | 868 | 44.7 | 5 | .3 | 233 | 151 | 366 | 750 | 229 | 135 | 364 | 18.7 | 3 10 | 5 19 1 | 63 | 3.2 | 183 | 152 |
| Sinoia | ... 173 | 1,109 | 764 | 2,196 | 587 | 2,267 | 1,351 | 4,463 | 188 | 16.9 | 80 | 7.2 | 115 | 10.4 | 34 | 3.0 | 42 | 3.8 | 24 | 2.2 | 176 | 15.9 | 404 | 36.4 | 4 | .4 | 168 | 102 | 219 | 489 | 140 | 72 | 212 | 19.1 | 3 10 | 5 3 8 | 42 | 3.8 | 105 | 108 |
| Belingwe | ... 60 | 509 | 365 | 730 | 50 | 1,865 | 415 | 2,595 | 69 | 13.6 | 42 | 8.2 | 38 | 7.4 | 7 | 1.4 | 9 | 1.8 | ... | ... | 69 | 13.6 | 246 | 48.3 | 8 | 1.6 | 50 | 14 | 226 | 290 | 59 | 59 | 11.6 | 3 5 | 7 10 0 | 21 | 4 1 | 79 | 105 | |
| Ingutsheni | ... 244 | 6,420* | 3,291 | 6,222 | 10,608 | 59,578 | 13,899 | 65,800 | 928 | 14.5 | 102 | 1.6 | 914 | 14.4 | 263 | 4.1 | 55 | .9 | 3 | ... | 1,530 | 23.8 | 2,489 | 38.8 | 40 | .6 | 344 | 680 | 9,181 | 10,155 | 334 | 705 | 1,039 | 16.2 | 1 7 | 18 19 3† | 96 | 1.5 | 192 | 233 |
| Totals | ... 5,594 | 43,810 | 28,668 | 55,013 | 42,690 | 145,633 | 71,358 | 200,646 | 6,437 | 14.7 | 4,374 | 10.0 | 3,444 | 7.9 | 2,866 | 6.5 | 1,130 | 2.6 | 455 | 1.0 | 7,764 | 17.7 | 15,513 | 35.4 | 387 | .9 | 12,219 | 4,212 | 19,142 | 35,573 | 9,694 | 3,632 | 13,326 | 30.4 | 3 3 | 5 9 0 | 1,440 | 3.3 | 6,004 | 6,879 |

* Includes value of supplies from hospital farm and garden, £752.

† In calculating this figure, the value of supplies from hospital farm and garden is deducted from total expenditure.

TABLE 22.

Return of Government and pauper patients treated in Government hospitals during 1924.

| Name of hospital. | Number of free patients. | | | Total number of units treated. | | | Cost of maintenance. £ s. d. | Loss of revenue represented, reckoning 5s. a day for whites, and 2s. 6d. a day for natives, plus extras. | | | | |
|--------------------------------|--------------------------|---------|---------|--------------------------------|-------------------------|---------|---------------------------------|--|----|--------|----|---|
| | White. | Native. | Totals. | White. | Native and coloured. | Totals. | | | | | | |
| | | | | | | £ | s. | d. | | | | |
| Salisbury ... | 248 | 502 | 750 | 5,218 | 15,121 | 20,339 | 4,841 | 14 | 0 | 3,194 | 12 | 6 |
| Umtali ... | 64 | 170 | 234 | 1,257 | 3,992 | 5,249 | 1,228 | 5 | 4 | 813 | 5 | 0 |
| Gwelo ... | 36 | 271 | 307 | 2,950 | 8,437 | 11,387 | 1,865 | 3 | 10 | 1,792 | 2 | 6 |
| Fort Victoria ... | 28 | 85 | 113 | 559 | 2,898 | 3,457 | 633 | 3 | 0 | 502 | 0 | 0 |
| Enkeldoorn ... | 8 | 41 | 49 | 166 | 813 | 979 | 243 | 3 | 8 | 143 | 2 | 6 |
| Gwanda ... | 20 | 50 | 70 | 114 | 1,102 | 1,216 | 191 | 0 | 8 | 166 | 5 | 0 |
| Gatooma ... | 21 | 887 | 908 | 242 | 20,231 | 20,473 | 2,400 | 9 | 2 | 2,589 | 7 | 6 |
| Shamva ... | 4 | 111 | 115 | 40 | 2,852 | 2,892 | 547 | 17 | 10 | 366 | 10 | 0 |
| Sinolia ... | 6 | 65 | 71 | 91 | 1,567 | 1,658 | 316 | 3 | 8 | 218 | 12 | 6 |
| Belingwe ... | 2 | 41 | 43 | 26 | 1,755 | 1,781 | 301 | 3 | 4 | 225 | 17 | 6 |
| Ingutsheni Mental Hospital ... | 36 | 181 | 217 | 9,570 | 53,906 | 63,476 | 5,112 | 19 | 10 | 9,130 | 15 | 0 |
| Totals ... | ... 473 | 2,404 | 2,877 | 20,233 | 112,674 | 132,907 | 17,681 | 4 | 4 | 19,142 | 10 | 0 |

